

ON THE FUNDAMENTAL GROUP OF THE COMPLEMENT OF TWO TANGENT CONICS AND AN ARBITRARY NUMBER OF TANGENT LINES

MEIRAV AMRAM^{1,2}, DAVID GARBER² AND MINA TEICHER

ABSTRACT. We compute the fundamental groups of the complements of the family of real conic-line arrangements with up to two conics which are tangent to each other at two points, with an arbitrary number of tangent lines to both conics. All the resulting groups turn out to be big.

Keywords: Conic-line arrangement, braid monodromy, fundamental group, tangent.

MSC 2000: Primary: 14H30; Secondary: 14J14, 14J17, 14Q05.

1. INTRODUCTION

Line arrangements as simply as they are, carry many open questions around them (including topological and combinatorial questions), which are slowly solved. In this paper, we go up to conic-line arrangements, where the parallel questions about them are even less understood.

We prove here a general theorem on the fundamental group's structure of complements of families of conic-line arrangements. The only known results so far in this direction are [4] and [5].

In general, the fundamental group of complements of plane curves is an important topological invariant with many different applications. Unfortunately, it is hard to compute.

¹Partially supported by the Emmy Noether Institute Fellowship (by the Minerva Foundation of Germany).

²Partially supported by a grant from the Ministry of Science, Culture and Sport, Israel and the Russian Foundation for Basic research, the Russian Federation.

Date: February 2, 2008.

This invariant was used by Chisini [8], Kulikov [13] and Kulikov-Teicher [14] in order to distinguish between different connected components of the moduli space of surfaces of general type.

Moreover, Zariski-Lefschetz hyperplane section theorem (see [15]) stated that

$$\pi_1(\mathbb{P}^N - S) \cong \pi_1(H - H \cap S),$$

where S is an hypersurface and H is a generic 2-plane. Since $H \cap S$ is a plane curve, we can compute $\pi_1(\mathbb{P}^N - S)$ in an easier way, by computing the fundamental group of the complement of a plane curve.

A different direction for the need of fundamental groups' computations is for getting more examples of Zariski pairs ([26],[27]). A pair of plane curves is called a *Zariski pair* if the two curves have the same combinatorics (i.e. the same singular points and the same arrangement of irreducible components), but their complements are not homeomorphic. Some examples of Zariski pairs can be found at [6], [7], [9], [10], [21], [22], [23] and [24].


Let $T_{n,m}$ be the family of real conic-line arrangements in $\mathbb{CP}^2$ with up to two conics, which are tangent to each other at two points, and with an arbitrary number of tangent lines to each one of the conics. The main result of this paper is the computation of the fundamental group $\pi_1(\mathbb{CP}^2 - T_{n,m})$:

Theorem 1.1. *Let Q_1, Q_2 be two tangent conics in $\mathbb{CP}^2$ and let $\{L_i\}_{i=1}^n$ and $\{L'_j\}_{j=1}^m$ be n and m lines which are tangent to Q_2 and Q_1 respectively, see Figure 1. Denote:*

$$T_{n,m} = Q_1 \cup Q_2 \cup \left(\bigcup_{i=1}^n L_i \right) \cup \left(\bigcup_{j=1}^m L'_j \right).$$

Then the fundamental group $\pi_1(\mathbb{CP}^2 - T_{n,m})$ is generated by the generators x_2 (related to Q_1), x_5 (related to Q_2), $x_6, \dots, x_{n+4}$ (related to $L_2, \dots, L_n$), $x_{n+5}, \dots, x_{n+m+4}$ (related to $L'_1, \dots, L'_m$). This group admits the following relations:

- (1) $x_{n+5}^{-1} x_5 x_{n+5} = x_{n+6}^{-1} \cdots x_{n+m+4}^{-1} \cdot x_5 \cdot x_{n+m+4} \cdots x_{n+6}$
- (2) $(x_2 x_i)^2 = (x_i x_2)^2$, where $i = 5, 6, \dots, n+4$
- (3) $(x_5 x_i)^2 = (x_i x_5)^2$, where $n+5 \leq i \leq n+m+4$


 FIGURE 1. The arrangement $T_{n,m}$

- (4) $[x_i, x_j] = e$, where $6 \leq i \leq n+4$ and $j = 5, n+5, \dots, n+m+4$
- (5) $[x_i, x_2x_5x_2^{-1}] = e$, where $6 \leq i \leq n+4$
- (6) $[x_2^{-1}x_ix_2, x_j] = e$, where $6 \leq i < j \leq n+4$
- (7) $[x_2, x_i] = e$, where $n+6 \leq i \leq n+m+4$
- (8) $[x_5x_2x_5^{-1}, x_i] = e$, where $n+5 \leq i \leq n+m+4$
- (9) $[x_{n+5}, x_i] = e$, where $i = 2, n+6, \dots, n+m+4$
- (10) $[x_5^{-1}x_ix_5, x_j] = e$, where $n+6 \leq i < j \leq n+m+4$.

We exclude here the case where at least one line is tangent to both conics (in their tangency point).

These arrangements may appear as a branch curve of a generic projection to $\mathbb{CP}^2$ of a surface of general type (see for example [12]).

A group is called *big* if it contains a subgroup, generated by two or more generators, which is free. By the above computations, we have the following corollary:

Corollary 1.2. *The fundamental group $\pi_1(\mathbb{CP}^2 - T_{n,m})$ is big.*

Algorithmically, this paper uses local computations (local braid monodromies and their induced relations, see also [1]), braid monodromy techniques of Moishezon-Teicher (see [11], [16], [17], [18], [19] and [20]), the van Kampen Theorem (see [25]) and some group simplifications and calculations for studying the fundamental groups (see also [2] and [3]).

Note that the new tool we are using in this paper is to construct the BMF of the general curve in an inductive way.

The paper is organized as follows. Section 2 deals with the fundamental group of a family of curves consisting of a conic and n tangent lines. In Section 3, we deal with the fundamental group of a family of curves consisting of two tangent conics and n tangent lines to one of the conics. In Section 4, we compute the most general case, where we have two tangent conics and two sets of lines, each of which is tangent to one of the conics.

In each section, we start with the computation of the braid monodromy of the curves by Moishezon-Teicher techniques and then we compute the simplified presentation of the fundamental groups by the van Kampen Theorem and group theoretic calculations.

2. THE FUNDAMENTAL GROUP OF THE COMPLEMENT OF A CONIC WITH n TANGENT LINES

In this section, we compute the fundamental group of the complement of a conic and n tangent lines. We start by illustrating the cases of one and two lines which are tangent to a conic, and then we present the computation of the general case of a conic and n tangent lines.

Note that this family of conic-line arrangements was already computed in [5] too (the family is denoted there by A_i). Although the presentations given here are different from those appeared in [5], it can be easily shown that the two presentations yield isomorphic groups.

We recompute their presentations here (by different methods) for the sake of completeness of presentations of conic-line arrangements with up to two tangent conics. One more reason for this is that it illustrates the method of computing the braid monodromy factorizations in an inductive way, as we will do in the case of two tangent conics and any number of tangent lines.

In the first subsection, we present the braid monodromy factorizations of this type of curves and in the next subsection, we compute the corresponding fundamental groups.

2.1. Braid monodromy factorizations. Here we compute the braid monodromy factorizations (BMFs, see [14]) which correspond to a conic with an arbitrary number of lines tangent to it. We start with some notations.

Notation 2.1. Let z_{ij} (resp. $\bar{z}_{ij}$) be a path below (resp. above) the axis, which connects points i and j . We denote by Z_{ij} (resp. $\bar{Z}_{ij}$) the counterclockwise half-twist of i and j along z_{ij} (resp. $\bar{z}_{ij}$).

We denote $Z_{i,jj'}^2 = Z_{ij'}^2 \cdot Z_{ij}^2$ and $Z_{i'i,jj'}^2 = Z_{i'j'}^2 \cdot Z_{i,jj'}^2$.

Conjugation of braids is denoted by $a^b = b^{-1}ab$.

Example 2.2. The path in Figure 2(a) is constructed as follows: take a path z_{34} and conjugate it by the full-twist $\bar{Z}_{13}^2$. Its corresponding half-twist is $Z_{34}^{\bar{Z}_{13}^2}$. The path in Figure 2(b) is constructed as follows: take again z_{34} and conjugate it first by Z_{23}^2 and then by Z_{13}^2 . Its corresponding half-twist is $Z_{34}^{Z_{23}^2 Z_{13}^2}$.



FIGURE 2. Examples of conjugated braids

Here, we start the computation of the BMFs.

Lemma 2.3. Let Q be a conic in $\mathbb{CP}^2$ and let L be a line which is tangent to Q (see Figure 3). Let $C_1 = Q \cup L$.




FIGURE 3. The arrangement C_1

Then its BMF is $\Delta_{C_1}^2 = F_1 \cdot (Z_{11'})^{Z_{1'2}^{-2}}$, where $F_1 = Z_{11'} \cdot (Z_{12}^4)^{Z_{11'}^2}$.

Proof. The computation of the BMF is done in a local model as follows: we start the computation by taking a typical fiber and enumerate the points of the curve in this fiber from 1 to n (n is the degree of the given curve). At this stage, we classify the singularities according to their types, and we summarize it in a table:

1	P_1	1	$\Delta^{\frac{1}{2}}_{\mathbb{R}I_2}\langle 1 \rangle$
2	$\langle 2, 3 \rangle$	4	$\Delta^2\langle 2, 3 \rangle$
3	$\langle 1, 2 \rangle$	1	$\Delta^{\frac{1}{2}}_{I_2\mathbb{R}}\langle 1 \rangle$

The second column contains the Lefschetz pairs corresponding to the singularities. In the third column, we indicate the types of the singularities, and in the last column we have the corresponding local diffeomorphisms of the singularities (see [3]).

In the next step, we apply the braid monodromy technique of Moishezon-Teicher [18] (see also [1] and [3]). In Figure 4, we show the skeletons of the braids, obtained by the technique. Note that we skip some steps in the braid monodromy's computation. Note also that at the end of the computation, we enumerate again the points in the fiber according to the global settings (for example, the two points of a conic are denoted by i and i').




FIGURE 4. The braid monodromy related to C_1

The first braid is $Z_{11'}$, the second one is $(Z_{12}^4)^{Z_{11'}^2}$ and the third one is $(Z_{11'})^{Z_{1'2}^{-2}}$, and hence the braid monodromy factorization is as stated.

■

Lemma 2.4. *Let Q be a conic in $\mathbb{CP}^2$ and let L_1, L_2 be two lines which are tangent to Q (see Figure 5). Let $C_2 = Q \cup L_1 \cup L_2$.*




FIGURE 5. The arrangement C_2

Then its BMF is $\Delta_{C_2}^2 = F_1 \cdot F_2 \cdot (Z_{11'})^{Z_{1'2}^{-2}Z_{1'3}^{-2}}$, where $F_1 = Z_{11'} \cdot (Z_{12}^4)^{Z_{11'}^2}$ and $F_2 = (Z_{23}^2)^{\bar{Z}_{12}^2} \cdot \bar{Z}_{13}^4$.

Proof. The table for computing the braid monodromy is:

1	P_1	1	$\Delta_{\mathbb{R}I_2}^{\frac{1}{2}} \langle 1 \rangle$
2	$\langle 2, 3 \rangle$	4	$\Delta^2 \langle 2, 3 \rangle$
3	$\langle 3, 4 \rangle$	2	$\Delta \langle 3, 4 \rangle$
4	$\langle 2, 3 \rangle$	4	$\Delta^2 \langle 2, 3 \rangle$
5	$\langle 1, 2 \rangle$	1	$\Delta_{I_2\mathbb{R}}^{\frac{1}{2}} \langle 1 \rangle$

In a similar way as in the previous lemma, we get the skeletons for the braid monodromies presented in Figure 6. ■

Theorem 2.5. *Let Q be a conic in $\mathbb{CP}^2$ and let $L_1, \dots, L_n$ be n lines which are tangent to Q . Let $C_n = Q \cup \bigcup_{i=1}^n L_i$ (see Figure 7).*

Then its BMF is:



$$\Delta_{C_n}^2 = F_1 \cdot F_2 \cdots F_n \cdot (Z_{11'})^{\prod_{i=1}^n Z_{1'i}^{-2}},$$

where:

$$F_1 = Z_{11'} \cdot (Z_{12}^4)^{Z_{11'}^2}$$

and

$$F_i = \prod_{k=2}^i (Z_{k,i+1}^2)^{\bar{Z}_{1k}^{-2}} \cdot \bar{Z}_{1,i+1}^4, \quad 2 \leq i \leq n.$$

FIGURE 6. The braid monodromy related to C_2 FIGURE 7. The arrangement C_n

Proof. We prove it by induction. The cases $n = 1, 2$ have been proved in the previous lemmas.

Assume that $\Delta_{C_n}^2$ is as given above. We will show that:

$$\Delta_{C_{n+1}}^2 = F_1 \cdot F_2 \cdots F_n \cdot F_{n+1} \cdot (Z_{11'})^{i=n+2} \prod_{i=n+2}^2 Z_{1',i}^{-2}.$$

The sub-factorization $F_1 \cdot F_2 \cdots F_n$ appears in $\Delta_{C_{n+1}}^2$, since by the structure of the configuration, the additional line L_{n+1} is located to the left of the rest of the lines $L_1, \dots, L_n$.

Note that by induction, adding the line L_{n+1} , causes no change in the factors F_i s, $1 \leq i \leq n$. This is due to the appearance of this line on the left side of $L_1, \dots, L_n$.

Now we explain the term:

$$F_{n+1} = \left(\prod_{k=2}^{n+1} (Z_{k,n+2}^2)^{\bar{Z}_{1k}^{-2}} \right) \cdot \bar{Z}_{1,n+2}^4.$$

The first part $\prod (Z_{k,n+2}^2)^{\bar{Z}_{1k}^{-2}}$ is the product of braids which correspond to the intersection points of L_{n+1} with the other lines $L_1, \dots, L_n$. Now, $\bar{Z}_{1,n+2}^4$ is the braid which corresponds to the tangency point of the line L_{n+1} and the conic Q .

We still have to explain the conjugated element. We have:

$$(Z_{11'})^{\prod_{i=n+2}^2 Z_{1',i}^{-2}} = \left((Z_{11'})^{\prod_{i=n+1}^2 Z_{1',i}^{-2}} \right)^{Z_{1',n+2}^{-2}},$$

where $(Z_{11'})^{\prod_{i=n+1}^2 Z_{1',i}^{-2}}$ appears in $\Delta_{C_n}^2$. Due to the addition of the line L_{n+1} , we have to conjugate this braid by the braid $Z_{1',n+2}^{-2}$. ■

2.2. The corresponding fundamental groups. Based on the BMFs we computed in the previous subsection, we now compute the fundamental groups of the affine and projective complements of the curve C_n .

Lemma 2.6. *Let C_1 be as in Lemma 2.3 (see Figure 3).*

Then we have:

$$\pi_1(\mathbb{C}^2 - C_1) \cong \mathbb{Z}^2 \quad ; \quad \pi_1(\mathbb{CP}^2 - C_1) \cong \mathbb{Z}.$$

Proof. By applying the van Kampen Theorem [25] on $\Delta_{C_1}^2$ (computed in Lemma 2.3), we get the following presentation for $\pi_1(\mathbb{C}^2 - C_1)$:

Generators: $\{x_1, x_{1'}, x_2\}$

Relations:

- (1) $x_1 = x_{1'}$
- (2) $(x_{1'} x_1 x_{1'}^{-1} x_2)^2 = (x_2 x_{1'} x_1 x_{1'}^{-1})^2$
- (3) $x_1 = x_{1'}^{-1} x_2^{-1} x_{1'} x_2 x_{1'}$

By Relation (1), the generator $x_{1'}$ is redundant, and by Relations (1) and (3), we get $x_1x_2 = x_2x_1$. Hence, we have:

$$\pi_1(\mathbb{C}^2 - C_1) = \langle x_1, x_2 \mid x_1x_2 = x_2x_1 \rangle \cong \mathbb{Z}^2.$$

Adding the projective relation $x_2x_{1'}x_1 = e$, we easily get:

$$\pi_1(\mathbb{CP}^2 - C_1) \cong \mathbb{Z}.$$

■

Lemma 2.7. *Let C_2 be as in Lemma 2.4 (see Figure 5).*

Then:

$$\pi_1(\mathbb{C}^2 - C_2) \cong \left\langle x_1, x_2, x_3 \left| \begin{array}{l} (x_1x_2)^2 = (x_2x_1)^2 \\ (x_1x_3)^2 = (x_3x_1)^2 \\ [x_3, x_1^{-1}x_2x_1] = [x_3x_2, x_1] = e \end{array} \right. \right\rangle,$$

where $[a, b] = aba^{-1}b^{-1}$ is the commutator element of a and b , and

$$\pi_1(\mathbb{CP}^2 - C_2) \cong \langle x_1, x_2 \mid (x_1x_2)^2 = (x_2x_1)^2 \rangle,$$

where the generator x_1 corresponds to the conic and the generators x_2, x_3 correspond to the lines.

Proof. By applying the van Kampen Theorem on $\Delta_{C_2}^2$ (computed in Lemma 2.4), we get the following presentation for $\pi_1(\mathbb{C}^2 - C_2)$:

Generators: $\{x_1, x_{1'}, x_2, x_3\}$

Relations:

- (1) $x_1 = x_{1'}$
- (2) $(x_{1'}x_1x_{1'}^{-1}x_2)^2 = (x_2x_{1'}x_1x_{1'}^{-1})^2$
- (3) $[x_3, x_2x_{1'}x_1x_{1'}^{-1}x_2x_{1'}x_1^{-1}x_{1'}^{-1}x_2^{-1}] = e$
- (4) $(x_2x_{1'}x_1x_{1'}^{-1}x_2^{-1}x_3)^2 = (x_3x_2x_{1'}x_1x_{1'}^{-1}x_2^{-1})^2$
- (5) $x_1 = x_{1'}^{-1}x_2^{-1}x_3^{-1}x_{1'}x_3x_2x_{1'}$

By Relation (1), the generator $x_{1'}$ is redundant. So we get:

Generators: $\{x_1, x_2, x_3\}$

Relations:

- (1) $(x_1x_2)^2 = (x_2x_1)^2$
- (2) $[x_3, x_2x_1x_2x_1^{-1}x_2^{-1}] = e$
- (3) $(x_2x_1x_2^{-1}x_3)^2 = (x_3x_2x_1x_2^{-1})^2$
- (4) $x_1 = x_2^{-1}x_3^{-1}x_1x_3x_2$

By Relation (1), Relation (2) is simplified to $[x_3, x_1^{-1}x_2x_1] = e$. Relation (4) is rewritten as $[x_1, x_3x_2] = e$. We treat now Relation (3):

$$x_2x_1x_2^{-1}x_3x_2x_1x_2^{-1}x_3 = x_3x_2x_1x_2^{-1}x_3x_2x_1x_2^{-1}.$$

By the relation $[x_3, x_1^{-1}x_2x_1] = e$, it can be written as:

$$x_2x_1x_2^{-1}x_1x_3x_3 = x_1x_3x_1x_3.$$

By Relation (4), $x_2x_1x_2^{-1} = x_3^{-1}x_1x_3$. So, we get $(x_1x_3)^2 = (x_3x_1)^2$.

The addition of the projective relation $x_3x_2x_1'x_1 = e$ enables us to omit the generator $x_3 = x_1^{-2}x_2^{-1}$. By substituting x_3 in any place it appears, we get the needed presentation for $\pi_1(\mathbb{CP}^2 - C_2)$. ■

Now, we compute the presentation for the general case.

Proposition 2.8. *Let Q be a conic in $\mathbb{CP}^2$ and let L_i ($1 \leq i \leq n$) be n lines which are tangent to Q (see Figure 7). Let $C_n = Q \cup \bigcup_{i=1}^n L_i$.*

Then:

$$\pi_1(\mathbb{C}^2 - C_n) \cong \left\langle x_1, x_2, \dots, x_{n+1} \left| \begin{array}{l} (x_1x_i)^2 = (x_ix_1)^2, \quad 2 \leq i \leq n+1 \\ [x_j, x_1^{-1}x_ix_1] = e, \quad 2 \leq i < j \leq n+1 \\ [x_{n+1} \cdots x_3x_2, x_1] = e \end{array} \right. \right\rangle$$

and

$$\pi_1(\mathbb{CP}^2 - C_n) \cong \left\langle x_1, x_2, \dots, x_{n+1} \left| \begin{array}{l} (x_1x_i)^2 = (x_ix_1)^2, \quad 2 \leq i \leq n+1 \\ [x_j, x_1^{-1}x_ix_1] = e, \quad 2 \leq i < j \leq n+1 \\ x_{n+1} \cdots x_3x_2x_1^2 = e \end{array} \right. \right\rangle,$$

where x_1 is a generator which corresponds to the conic and $x_2, \dots, x_{n+1}$ are generators which correspond to the n lines.

Proof. In this case, the set of generators for the presentation of the fundamental group is $\{x_1, x_1', x_2, \dots, x_{n+1}\}$. By applying the van Kampen Theorem on the braid monodromy computed in Theorem 2.5, we get three types of relations:

(1) From the branch points, we get $x_1 = x_1'$ and

$$x_1 = x_1^{-1}x_2^{-1}x_3^{-1} \cdots x_{n+1}^{-1}x_1'x_{n+1} \cdots x_3x_2x_1.$$

(2) From the nodal points, we get the following set of relations:

$$[x_j, x_{j-1}x_{j-2} \cdots x_{1'}x_1x_{1'}^{-1}x_2^{-1} \cdots x_{i-1}^{-1}x_ix_{i-1} \cdots x_2x_{1'}x_1^{-1}x_{1'}^{-1} \cdots x_{j-2}^{-1}x_{j-1}^{-1}] = e,$$

where $2 \leq i < j \leq n+1$.

(3) From the tangency points, we get:

$$(x_1x_2^{-1}x_3^{-1} \cdots x_{i-1}^{-1}x_ix_{i-1} \cdots x_3x_2)^2 = (x_2^{-1}x_3^{-1} \cdots x_{i-1}^{-1}x_ix_{i-1} \cdots x_3x_2x_1)^2,$$

where $2 \leq i \leq n+1$.

By the relations of the first type (from the branch points), $x_{1'}$ is redundant, and we have: $[x_{n+1} \cdots x_3x_2, x_1] = e$.

Now, we simplify the relations of types (2) and (3). In type (3), we get for $i = 2$: $(x_1x_2)^2 = (x_2x_1)^2$. In type (2), for $i = 2$ and $j = 3$, we get: $[x_3, x_2x_1x_2x_1^{-1}x_2^{-1}] = e$. By $(x_1x_2)^2 = (x_2x_1)^2$, it is $[x_3, x_1^{-1}x_2x_1] = e$.

Returning again to type (3), for $i = 3$, we have: $(x_1x_2^{-1}x_3x_2)^2 = (x_2^{-1}x_3x_2x_1)^2$, which is:

$$(x_1x_2^{-1}x_3x_2)(x_1x_2^{-1}x_3x_2) = (x_2^{-1}x_3x_2x_1)(x_2^{-1}x_3x_2x_1).$$

Since $(x_1x_2)^2 = (x_2x_1)^2$, we have

$$x_1x_2^{-1}x_3x_1^{-1}x_2^{-1}x_1x_2x_1x_3x_2 = x_2^{-1}x_3x_1^{-1}x_2^{-1}x_1x_2x_1x_3x_2x_1,$$

and by $[x_3, x_1^{-1}x_2x_1] = e$, we get:

$$x_1x_2^{-1}x_1^{-1}x_2^{-1}x_1x_3x_2x_1x_3x_2 = x_2^{-1}x_1^{-1}x_2^{-1}x_1x_3x_2x_1x_3x_2x_1.$$

This relation is simplified to (by $(x_1x_2)^2 = (x_2x_1)^2$):

$$x_1x_3x_2x_1x_3x_2 = x_3x_2x_1x_3x_2x_1.$$

This relation is:

$$x_1x_3x_2x_1x_3 = x_3x_2x_1x_3x_2x_1x_2^{-1}.$$

Using $(x_1x_2)^2 = (x_2x_1)^2$, we have:

$$x_1x_3x_2x_1x_3 = x_3x_2x_1x_3x_1^{-1}x_2^{-1}x_1x_2x_1.$$

Now we add $x_1x_1^{-1} = e$:

$$x_1x_3(x_1x_1^{-1})x_2x_1x_3 = x_3x_2x_1x_3x_1^{-1}x_2^{-1}x_1x_2x_1.$$

Since $[x_3, x_1^{-1}x_2x_1] = e$,

$$x_1x_3x_1x_3x_1^{-1}x_2x_1 = x_3x_2x_1x_1^{-1}x_2^{-1}x_1x_3x_2x_1.$$

And we get $(x_1x_3)^2 = (x_3x_1)^2$.

Now we return again to type (2), for $i = 2$ and $j = 4$, we have: $[x_4, x_3x_2x_1x_2x_1^{-1}x_2^{-1}x_3^{-1}] = e$. By $(x_1x_2)^2 = (x_2x_1)^2$, it is: $[x_4, x_3x_1^{-1}x_2x_1x_3^{-1}] = e$, which is simplified to: $[x_4, x_1^{-1}x_2x_1] = e$, by using $[x_3, x_1^{-1}x_2x_1] = e$.

Now, for $i = 3$ and $j = 4$, we have: $[x_4, x_3x_2x_1x_2^{-1}x_3x_2x_1^{-1}x_2^{-1}x_3^{-1}] = e$. By $(x_1x_2)^2 = (x_2x_1)^2$, we rewrite it as:

$$[x_4, x_3x_1^{-1}x_2^{-1}x_1x_2x_1x_3x_1^{-1}x_2^{-1}x_1^{-1}x_2x_1x_3^{-1}] = e,$$

which is (by $[x_3, x_1^{-1}x_2x_1] = e$):

$$[x_4, x_1^{-1}x_2^{-1}x_1x_3x_2x_1x_3x_1^{-1}x_2^{-1}x_3^{-1}x_1^{-1}x_2x_1] = e.$$

Now, by $[x_4, x_1^{-1}x_2x_1] = e$:

$$[x_4, x_3x_2x_1x_3x_1^{-1}x_2^{-1}x_3^{-1}] = e.$$

By $[x_3, x_1^{-1}x_2x_1] = e$, we get: $[x_4, x_3x_1x_3x_1^{-1}x_3^{-1}] = e$ and by $(x_1x_3)^2 = (x_3x_1)^2$, we get $[x_4, x_1^{-1}x_3x_1] = e$.

In the same way, we get the following sets of relations:

$$(1) \quad [x_j, x_1^{-1}x_ix_1] = e, \quad \text{where } 2 \leq i < j \leq n+1$$

and

$$(2) \quad (x_1x_i)^2 = (x_ix_1)^2 \quad \text{where } 2 \leq i \leq n+1,$$

as required.

Adding the projective relation $x_{n+1}x_n \cdots x_2x_1^2 = e$, the relation

$$[x_{n+1} \cdots x_3x_2, x_1] = e$$

is redundant, so we get the needed presentation for the projective case too. ■

2.3. The groups are big. Following the results of the previous subsections, we have:

Corollary 2.9. *The affine and projective fundamental groups of a conic-line arrangement, composed of one conic and at least two tangent lines, are big.*

We have already proved a similar corollary in [1], and for the sake of completeness, we recall its proof shortly here.

Proof. Observe that if a group has a big quotient, then the group itself is big.

Denote $G = \pi_1(\mathbb{CP}^2 - C_2) \cong \langle a, b \mid (ab)^2 = (ba)^2 \rangle$. We show that G is big. We start by taking the quotient by the relation $(ab)^2 = e$. Then we get the group $\langle a, b \mid (ab)^2 = (ba)^2 = e \rangle$.

Let us take new generators $x = ab, y = b$, then the relation $(ab)^2 = (ba)^2 = e$ becomes: $x^2 = (yxy^{-1})^2 = e$, which is equal to: $x^2 = yx^2y^{-1} = e$. Hence, we have:

$$\langle a, b \mid (ab)^2 = (ba)^2 = e \rangle \cong \langle x, y \mid x^2 = e \rangle \cong \mathbb{Z} * \mathbb{Z}/2,$$

where $*$ is the free product.

Now, the quotient of $G/\langle x^2 = e \rangle$ by the subgroup generated by y^3 is $\mathbb{Z}/2 * \mathbb{Z}/3$, which is known to be big. By the observation, $G = \pi_1(\mathbb{CP}^2 - C_2)$ is big too.

Since C_2 is a sub-arrangement of C_n , $n \geq 2$, the group $G = \pi_1(\mathbb{CP}^2 - C_2)$ is a quotient of the group $\pi_1(\mathbb{CP}^2 - C_n)$ (by sending the generators which correspond to the additional lines to e). Now, since G is big, the above groups are big as well.

The above proof shows that the projective fundamental group of C_n , $n \geq 2$, is big. Since the projective fundamental group is a quotient of the affine fundamental group (by the projective relation), the corresponding affine fundamental groups are also big, again by the observation, as needed. ■

3. THE FUNDAMENTAL GROUP OF THE COMPLEMENT OF TWO TANGENT CONICS WITH n TANGENT LINES TO THE SAME CONIC

In this section, we compute the fundamental group of the projective complement of a curve composed of two tangent conics and n additional lines, which are tangent to the same conic.

As before, in the first subsection, we present the BMF of this type of curves and in the second subsection, we compute the corresponding fundamental groups.

3.1. The braid monodromy factorizations. Let $T_{n,0}$ be a conic-line arrangement, composed of two tangent conics (which are tangent to each other at two points), and with n additional lines (which are tangent to the same conic).

We start with the arrangements $T_{0,0}$, $T_{1,0}$ and $T_{2,0}$. Their fundamental groups have been already computed in [1].

We give here their BMFs:

Lemma 3.1. *Let $T_{0,0}$ be a conic arrangement composed of two conics which are tangent at two points (see Figure 8).*





FIGURE 8. The arrangement $T_{0,0}$

The BMF of $T_{0,0}$ is:

$$\Delta_{T_{0,0}}^2 = Z_{34} \cdot Z_{24}^4 \cdot (Z_{12})^{Z_{24}^2} \cdot (Z_{12})^{Z_{23}^2} \cdot Z_{23}^4 \cdot Z_{34}.$$


Lemma 3.2. *Let $T_{1,0}$ be a conic-line arrangement composed of two conics which are tangent at two points, and a line which is tangent to one of the conics (see Figure 9).*

FIGURE 9. The arrangement $T_{1,0}$

Then its BMF is:

$$\begin{aligned} \Delta_{T_{1,0}}^2 = & Z_{45} \cdot Z_{12}^2 \cdot Z_{35}^4 \cdot (Z_{13}^2)^{Z_{35}^2 Z_{12}^2} \cdot (Z_{23})^{Z_{35}^2} \cdot (Z_{23})^{Z_{12}^{-2} Z_{34}^2} \cdot \\ & \cdot (Z_{15}^4)^{Z_{12}^2} \cdot Z_{34}^4 \cdot (Z_{45})^{Z_{14}^{-2} Z_{12}^2}. \end{aligned}$$


Lemma 3.3. Let $T_{2,0}$ be a conic-line arrangement composed of two conics which are tangent at two points, and two lines which are tangent to the same conic (see Figure 10).

FIGURE 10. The arrangement $T_{2,0}$

Then its BMF is:

$$\begin{aligned} \Delta_{T_{2,0}}^2 = & Z_{23}^2 \cdot Z_{56} \cdot Z_{46}^4 \cdot Z_{23}^2 \cdot (Z_{34})^{Z_{23}^2 Z_{46}^2} \cdot (Z_{34})^{Z_{23}^{-2} Z_{45}^2} \cdot Z_{45}^4 \cdot (Z_{26}^4)^{Z_{23}^2} \cdot \\ & (Z_{12}^2)^{Z_{26}^2 Z_{23}^2} \cdot Z_{16}^4 \cdot (Z_{56})^{Z_{15}^{-2} Z_{25}^{-2} Z_{23}^2} \cdot (Z_{14}^2)^{Z_{12}^2 Z_{23}^2} Z_{13}^2. \end{aligned}$$

Now, let us consider the conic-line arrangement $T_{n,0}$, which is composed of two tangent conics (which are tangent to each other at two points) and n lines which are tangent to the same conic.

FIGURE 11. The arrangement $T_{n,0}$

Proposition 3.4. *Let Q_1, Q_2 be two tangent conics in $\mathbb{CP}^2$ (which are tangent to each other at two points) and let $L_1, \dots, L_n$ be n lines tangent to Q_2 (see Figure 11). Denote*

$$T_{n,0} = Q_1 \cup Q_2 \cup \bigcup_{i=1}^n L_i.$$

Then its BMF is:

$$\begin{aligned} \Delta_{T_{n,0}}^2 &= Z_{n+2,n+3} \cdot (Z_{n,n+1})^{Z_{n-1,n}^2 Z_{n+1,n+3}^2} \cdot (Z_{n,n+1})^{Z_{n-1,n}^2 Z_{n+1,n+2}^{-2}} \cdot \\ &\quad \cdot \tilde{Z}_{n+2,n+3} \cdot Z_{n+1,n+3}^4 \cdot Z_{n+1,n+2}^4 \cdot (Z_{n-1,n+3}^4)^{Z_{n-1,n}^2} \cdot Z_{n+2,n+4}^4 \cdot \\ &\quad \cdot \prod_{i=1}^{n-2} Z_{i,n+3}^4 \cdot \prod_{i=1}^{n-2} (Z_{i,n-1}^2)^{Z_{n-1,n+3}^2 Z_{n-1,n}^2} \cdot \prod_{i=1}^{n-2} \tilde{Z}_{i,n+4}^2 \cdot \\ &\quad \cdot \prod_{1 \leq i < j \leq n-2} (Z_{i,j}^2)^{Z_{j,n+3}^2} \cdot \prod_{i=1}^{n-2} \tilde{Z}_{i,n}^2 \cdot \prod_{i=1}^{n-2} \tilde{Z}_{i,n+1}^2 \cdot Z_{n-1,n}^2 \cdot \\ &\quad \cdot (\bar{Z}_{n,n+4}^2)^{Z_{n-1,n}^2} \cdot \tilde{Z}_{n+1,n+4}^2 \cdot (Z_{n-1,n+4}^2)^{Z_{n-1,n}^2 Z_{n+3,n+4}^{-2}} \end{aligned}$$

The skeletons of the braids $\tilde{Z}_{n+2,n+3}$, $\tilde{Z}_{i,n+4}^2$, $\tilde{Z}_{i,n}^2$, $\tilde{Z}_{i,n+1}^2$ and $\tilde{Z}_{n+1,n+4}^2$ appear in Figure 12.




FIGURE 12. Skeletons for some braids appearing in the BMF of $T_{n,0}$

Before the proof, we have the following remark.

Remark 3.5. *The order of the factors in a BMF Δ^2 is fixed according to the order of the locations of the singularities in the curve (see [20]). Since our goal is to compute fundamental groups, the order of the factors is irrelevant. Hence, we list here the monodromy factors without preserving their original order (we join together sets of monodromies of the same type), and concentrate on finding the relations in the group by applying the van Kampen Theorem on the monodromies (a similar convention was also applied in [2], since we were also interested in the presentation and not in the actual order in the factorization).*

Proof of Proposition 3.4. The first four factors in $\Delta_{T_{n,0}}^2$ correspond to the four branch points in the arrangement.

The next five factors correspond to the tangent points in the arrangement (two of them correspond to the tangency points between the two

conics, and the others correspond to the tangency points between one of the conics and the lines).

The last nine factors correspond to the intersections of the lines with the second conic, and between the lines themselves. $\blacksquare$

3.2. The corresponding fundamental groups. In this section, we compute the projective fundamental groups $\pi_1(\mathbb{CP}^2 - T_{n,0})$, using the BMFs we computed in the previous section (as in [1]). The fundamental groups of the complements of $T_{0,0}$, $T_{1,0}$ and $T_{2,0}$ have been computed in [1] (Propositions 1.1, 1.2, 1.3), therefore we quote here the results.

Proposition 3.6. *Let $T_{0,0}$ be the curve defined in Lemma 3.1 (see Figure 8). Then:*

$$\pi_1(\mathbb{CP}^2 - T_{0,0}) \cong \langle x_1, x_2 \mid (x_1x_2)^2 = (x_2x_1)^2 = e \rangle \cong \mathbb{Z} * \mathbb{Z}/2.$$

Proposition 3.7. *Let $T_{1,0}$ be the curve defined in Lemma 3.2 (see Figure 9). Then:*

$$\pi_1(\mathbb{CP}^2 - T_{1,0}) \cong \langle x_1, x_2 \mid (x_1x_2)^2 = (x_2x_1)^2 \rangle.$$

Proposition 3.8. *Let $T_{2,0}$ be the curve defined in Lemma 3.3 (see Figure 10). Then:*

$$\pi_1(\mathbb{CP}^2 - T_{2,0}) \cong \left\langle x_1, x_2, x_3 \mid \begin{array}{l} (x_2x_3)^2 = (x_3x_2)^2, (x_1x_3)^2 = (x_3x_1)^2, \\ [x_1, x_2] = [x_2, x_3x_1x_3^{-1}] = e \end{array} \right\rangle.$$

Now we proceed to the general case:

Theorem 3.9. *Let Q_1, Q_2 be two tangent conics in $\mathbb{CP}^2$ (which are tangent to each other at two points) and let $L_1, \dots, L_n$ be n lines tangent to Q_2 (see Figure 11). Denote*

$$T_{n,0} = Q_1 \cup Q_2 \cup \bigcup_{i=1}^n L_i,$$

as in Proposition 3.4. Then:

$$\pi_1(\mathbb{CP}^2 - T_{n,0}) \cong \left\langle x_1, \dots, x_n, x_{n+2} \mid \begin{array}{ll} (x_nx_{n+2})^2 = (x_{n+2}x_n)^2 & \\ (x_ix_{n+2})^2 = (x_{n+2}x_i)^2, & 1 \leq i \leq n-1 \\ [x_i, x_n] = [x_i, x_{n+2}x_nx_{n+2}^{-1}] = e, & 1 \leq i \leq n-1 \\ [x_i, x_{n+2}x_jx_{n+2}^{-1}] = e, & 1 \leq i < j \leq n-1 \end{array} \right\rangle.$$

The generators $x_1, \dots, x_{n-1}$ correspond to the lines, and the generators x_n and x_{n+2} correspond to the two conics.

Proof. Applying the van Kampen Theorem on $\Delta_{T_{n,0}}^2$ (computed in Proposition 3.4), we get the following presentation for $\pi_1(\mathbb{CP}^2 - T_{n,0})$:

Generators: $\{x_1, x_2, \dots, x_{n+4}\}$

Relations:

- (1) $x_{n+2} = x_{n+3}$
- (2) $x_n x_{n-1} x_n x_{n-1}^{-1} x_n^{-1} = x_{n+3} x_{n+1} x_{n+3}^{-1}$
- (3) $x_n x_{n-1} x_n x_{n-1}^{-1} x_n^{-1} = x_{n+1}^{-1} x_{n+2}^{-1} x_{n+1} x_{n+2} x_{n+1}$
- (4) $x_{n+3} = x_1^{-1} x_2^{-1} \cdots x_{n-2}^{-1} x_n x_{n-1}^{-1} x_n^{-1} x_{n+3} x_{n+4} x_{n+3} x_{n+2} x_{n+3}^{-1} x_{n+4}^{-1} x_{n+3} x_n x_{n-1} x_n^{-1} x_{n-2} \cdots x_2 x_1$
- (5) $(x_{n+1} x_{n+3})^2 = (x_{n+3} x_{n+1})^2$
- (6) $(x_{n+1} x_{n+2})^2 = (x_{n+2} x_{n+1})^2$
- (7) $(x_n x_{n-1} x_n^{-1} x_{n+3})^2 = (x_{n+3} x_n x_{n-1} x_n^{-1})^2$
- (8) $(x_{n+2} x_{n+4})^2 = (x_{n+4} x_{n+2})^2$
- (9) $(x_i x_{n+3})^2 = (x_{n+3} x_i)^2$, where $1 \leq i \leq n-2$
- (10) $[x_i, x_{n+3} x_n x_{n-1} x_n^{-1} x_{n+3}] = e$, where $1 \leq i \leq n-2$
- (11) $[x_{n+3} x_{n+4} x_{n+3}, x_{n-1}^{-1} x_n x_{n-1} x_{n-2} \cdots x_{i+1} x_i x_{i+1}^{-1} \cdots x_{n-2}^{-1} x_{n-1}^{-1} x_{n-1}] = e$, where $1 \leq i \leq n-2$
- (12) $[x_i, x_{n+3} x_j x_{n+3}^{-1}] = e$, where $1 \leq i < j \leq n-2$
- (13) $[x_n, x_{n-2} \cdots x_{i+1} x_i x_{i+1}^{-1} \cdots x_{n-2}^{-1}] = e$ where $1 \leq i \leq n-2$.
- (14) $[x_n x_{n-1}^{-1} x_n^{-1} x_{n+1} x_n x_{n-1} x_n^{-1}, x_{n-2} \cdots x_{i+1} x_i x_{i+1}^{-1} \cdots x_{n-2}^{-1}] = e$, where $1 \leq i \leq n-2$
- (15) $[x_{n-1}, x_n] = e$
- (16) $[x_n x_{n-1} x_n x_{n-1}^{-1} x_n^{-1}, x_{n+1}^{-1} x_{n+2}^{-1} x_{n+3}^{-1} x_{n+4} x_{n+3} x_{n+2} x_{n+1}] = e$
- (17) $[x_{n+1}, x_{n+2} x_{n+4} x_{n+2}^{-1} x_{n+4} x_{n+2} x_{n+4} x_{n+2}] = e$
- (18) $[x_n x_{n-1} x_n^{-1}, x_{n+3} x_{n+4} x_{n+3}] = e$
- (19) $x_{n+4} x_{n+3} x_{n+2} x_{n+1} x_n \cdots x_1 = e$ (Projective relation).

By Relation (1), the generator x_{n+3} can be omitted, and we replace it with x_{n+2} in any place it appears. Moreover, using Relation (15), we can simplify many relations, and we get the following presentation.

Generators: $\{x_1, \dots, x_{n+2}, x_{n+4}\}$

Relations:

- (1) $x_n = x_{n+2} x_{n+1} x_{n+2}^{-1}$
- (2) $x_n = x_{n+1}^{-1} x_{n+2}^{-1} x_{n+1} x_{n+2} x_{n+1}$
- (3) $x_{n+2} = x_1^{-1} x_2^{-1} \cdots x_{n-2}^{-1} x_n x_{n-1}^{-1} x_n^{-1} x_{n+2} x_{n+4} x_{n+2} x_{n+4}^{-1} x_{n+2} x_{n-1} x_{n-2} \cdots x_2 x_1$
- (4) $(x_{n+1} x_{n+2})^2 = (x_{n+2} x_{n+1})^2$
- (5) $(x_{n-1} x_{n+2})^2 = (x_{n+2} x_{n-1})^2$
- (6) $(x_{n+2} x_{n+4})^2 = (x_{n+4} x_{n+2})^2$
- (7) $(x_i x_{n+2})^2 = (x_{n+2} x_i)^2$, where $1 \leq i \leq n-2$
- (8) $[x_{n-1}, x_n] = e$
- (9) $[x_i, x_{n+2} x_{n-1} x_{n+2}^{-1}] = e$, where $1 \leq i \leq n-2$
- (10) $[x_{n+2} x_{n+4} x_{n+2}, x_n x_{n-2} \cdots x_{i+1} x_i x_{i+1}^{-1} \cdots x_{n-2}^{-1} x_n^{-1}] = e$, where $1 \leq i \leq n-2$
- (11) $[x_i, x_{n+2} x_j x_{n+2}^{-1}] = e$, where $1 \leq i < j \leq n-2$
- (12) $[x_n, x_{n-2} \cdots x_{i+1} x_i x_{i+1}^{-1} \cdots x_{n-2}^{-1}] = e$, where $1 \leq i \leq n-2$

- (13) $[x_{n-1}^{-1}x_{n+1}x_{n-1}, x_{n-2} \cdots x_{i+1}x_ix_{i+1}^{-1} \cdots x_{n-2}^{-1}] = e$, where $1 \leq i \leq n-2$
- (14) $[x_n, x_{n+1}^{-1}x_{n+2}^{-2}x_{n+4}x_{n+2}^2x_{n+1}] = e$
- (15) $[x_{n+1}, x_{n+2}^{-1}x_{n+4}^{-1}x_{n+2}^{-1}x_{n+4}x_{n+2}x_{n+4}x_{n+2}] = e$
- (16) $[x_{n-1}, x_{n+2}^{-1}x_{n+4}x_{n+2}] = e$
- (17) $x_{n+4}x_{n+2}^2x_{n+1}x_n \cdots x_1 = e$.

By Relation (6), Relation (15) is simplified to $[x_{n+1}, x_{n+4}] = e$. We combine Relations (4)–(7) together, and Relations (9) and (11) together. So, we get the following set of relations:

- (1) $x_n = x_{n+2}x_{n+1}x_{n+2}^{-1}$
- (2) $x_n = x_{n+1}^{-1}x_{n+2}^{-1}x_{n+1}x_{n+2}x_{n+1}$
- (3) $x_{n+2} = x_1^{-1}x_2^{-1} \cdots x_{n-2}^{-1}x_{n-1}^{-1}x_{n+2}^{-1}x_{n+4}x_{n+2}x_{n+4}^{-1}x_{n+2}x_{n-1}x_{n-2} \cdots x_2x_1$
- (4) $(x_ix_{n+2})^2 = (x_{n+2}x_i)^2$, where $i = 1, \dots, n-1, n+1, n+4$
- (5) $[x_{n-1}, x_n] = e$
- (6) $[x_{n+2}^{-1}x_{n+4}x_{n+2}, x_nx_{n-2} \cdots x_{i+1}x_ix_{i+1}^{-1} \cdots x_{n-2}^{-1}x_n^{-1}] = e$, where $1 \leq i \leq n-2$
- (7) $[x_i, x_{n+2}x_jx_{n+2}^{-1}] = e$, where $1 \leq i < j \leq n-1$
- (8) $[x_n, x_{n-2} \cdots x_{i+1}x_ix_{i+1}^{-1} \cdots x_{n-2}^{-1}] = e$, where $1 \leq i \leq n-2$
- (9) $[x_{n-1}^{-1}x_{n+1}x_{n-1}, x_{n-2} \cdots x_{i+1}x_ix_{i+1}^{-1} \cdots x_{n-2}^{-1}] = e$, where $1 \leq i \leq n-2$
- (10) $[x_n, x_{n+1}^{-1}x_{n+2}^{-2}x_{n+4}x_{n+2}^2x_{n+1}] = e$
- (11) $[x_{n+1}, x_{n+4}] = e$
- (12) $[x_{n-1}, x_{n+2}^{-1}x_{n+4}x_{n+2}] = e$
- (13) $x_{n+4}x_{n+2}^2x_{n+1}x_n \cdots x_1 = e$.

We treat Relations (8). For $i = n-2$, we get $[x_n, x_{n-2}] = e$. Then, for $i = n-3$, we get $[x_n, x_{n-2}x_{n-3}x_{n-2}^{-1}] = e$. By the relation $[x_n, x_{n-2}] = e$, we can simplify it to $[x_n, x_{n-3}] = e$. If we continue in a decreasing order of the indices $i = n-4, \dots, 1$, we get: $[x_n, x_i] = e$ for $1 \leq i \leq n-2$. Now we can combine this set with Relation (5).

In a similar way, we can simplify Relations (9) to get for $1 \leq i \leq n-2$:

$$[x_{n-1}^{-1}x_{n+1}x_{n-1}, x_i] = e.$$

This simplification gives us the following list of relations:

- (1) $x_n = x_{n+2}x_{n+1}x_{n+2}^{-1}$
- (2) $x_n = x_{n+1}^{-1}x_{n+2}^{-1}x_{n+1}x_{n+2}x_{n+1}$
- (3) $x_{n+2} = x_1^{-1}x_2^{-1} \cdots x_{n-2}^{-1}x_{n-1}^{-1}x_{n+2}^{-1}x_{n+4}x_{n+2}x_{n+4}^{-1}x_{n+2}x_{n-1}x_{n-2} \cdots x_2x_1$
- (4) $(x_ix_{n+2})^2 = (x_{n+2}x_i)^2$, where $i = 1, \dots, n-1, n+1, n+4$
- (5) $[x_i, x_n] = e$, where $1 \leq i \leq n-1$
- (6) $[x_{n+2}^{-1}x_{n+4}x_{n+2}, x_nx_{n-2} \cdots x_{i+1}x_ix_{i+1}^{-1} \cdots x_{n-2}^{-1}x_n^{-1}] = e$, where $1 \leq i \leq n-2$
- (7) $[x_i, x_{n+2}x_jx_{n+2}^{-1}] = e$, where $1 \leq i < j \leq n-1$
- (8) $[x_{n-1}^{-1}x_{n+1}x_{n-1}, x_i] = e$, where $1 \leq i \leq n-2$
- (9) $[x_n, x_{n+1}^{-1}x_{n+2}^{-2}x_{n+4}x_{n+2}^2x_{n+1}] = e$
- (10) $[x_{n+1}, x_{n+4}] = e$
- (11) $[x_{n-1}, x_{n+2}^{-1}x_{n+4}x_{n+2}] = e$
- (12) $x_{n+4}x_{n+2}^2x_{n+1}x_n \cdots x_1 = e$.

By substituting Relation (1) in Relations (4) for $i = n + 1$, we get $(x_n x_{n+2})^2 = (x_{n+2} x_n)^2$. Since the relation $(x_{n+1} x_{n+2})^2 = (x_{n+2} x_{n+1})^2$ is known in Relations (4), Relation (2) is redundant (since comparing Relations (1) and (2) yields a known relation). Now we substitute Relation (1) in Relation (9) and we get:

$$\begin{aligned} e &= [x_{n+1} x_n x_{n+1}^{-1}, x_{n+2}^{-2} x_{n+4} x_{n+2}^2] = [(x_{n+2}^{-1} x_n x_{n+2}) x_n (x_{n+2}^{-1} x_n^{-1} x_{n+2}), x_{n+2}^{-2} x_{n+4} x_{n+2}^2] = \\ &= [x_n x_{n+2} x_n x_{n+2}^{-1}, x_{n+2}^{-1} x_{n+4} x_{n+2}] = [x_{n+2}^{-1} x_n x_{n+2}, x_{n+2}^{-1} x_{n+4} x_{n+2}] = [x_n, x_{n+4}]. \end{aligned}$$

By Relation (12), we substitute $x_{n+2}^2 x_{n+1} x_n \cdots x_1 = x_{n+4}^{-1}$ in $[x_n, x_{n+4}] = e$. By Relations (1) and (5), this relation can be simplified to $(x_{n+2} x_n)^2 = (x_n x_{n+2})^2$, which is known, and hence Relation (9) is redundant. So we get the following set of relations:

- (1) $x_n = x_{n+2} x_{n+1} x_{n+2}^{-1}$
- (2) $x_{n+2} = x_1^{-1} x_2^{-1} \cdots x_{n-2}^{-1} x_{n-1}^{-1} x_{n+2} x_{n+4} x_{n+2}^{-1} x_{n+4} x_{n+2} x_{n-1} x_{n-2} \cdots x_2 x_1$
- (3) $(x_i x_{n+2})^2 = (x_{n+2} x_i)^2$, where $i = 1, \dots, n+1, n+4$
- (4) $[x_i, x_n] = e$, where $1 \leq i \leq n-1$
- (5) $[x_{n+2}^{-1} x_{n+4} x_{n+2}, x_n x_{n-2} \cdots x_{i+1} x_i x_{i+1}^{-1} \cdots x_{n-2}^{-1} x_n^{-1}] = e$, where $1 \leq i \leq n-2$
- (6) $[x_i, x_{n+2} x_j x_{n+2}^{-1}] = e$, where $1 \leq i < j \leq n-1$
- (7) $[x_{n-1}^{-1} x_{n+1} x_{n-1}, x_i] = e$, where $1 \leq i \leq n-2$
- (8) $[x_{n+1}, x_{n+4}] = e$
- (9) $[x_{n-1}, x_{n+2}^{-1} x_{n+4} x_{n+2}] = e$
- (10) $x_{n+4} (x_{n+2} x_n)^2 x_{n-1} \cdots x_1 = e$.

We start by simplifying Relations (5) for $i = n-2$:

$$e = [x_{n+2}^{-1} x_{n+4} x_{n+2}, x_n x_{n-2} x_n^{-1}] \stackrel{(4), i=n-2}{=} [x_{n+4}, x_{n+2} x_{n-2} x_{n+2}^{-1}].$$

This relation can be rewritten, by using Relation (10), as:

$$x_n x_{n+2} x_n x_{n-1} \cdots x_1 x_{n+2} x_{n-2} x_{n+2}^{-1} = x_{n-2} x_n x_{n+2} x_n x_{n-1} \cdots x_1.$$

Now, by Relations (3) (for $i = n-2$), we get:

$$x_n x_{n+2} x_n x_{n-1} x_{n-2} x_{n+2} x_{n-2} x_{n+2}^{-1} x_{n-2}^{-1} = x_{n-2} x_n x_{n+2} x_n x_{n-1}.$$

By Relations (6) (for $i = n-2$), we get:

$$x_{n+2} x_n x_{n+2}^{-1} x_{n-2} = x_{n-2} x_{n+2} x_n x_{n+2}^{-1}.$$

In a similar way, we get $[x_i, x_{n+2} x_n x_{n+2}^{-1}] = e$, for $1 \leq i \leq n-3$. As done above, we substitute this time Relation (10) in Relation (9) to

get: $[x_{n-1}, x_{n+2}x_nx_{n+2}^{-1}] = e$. We combine both induced relations in Relations (5) below, to get:

- (1) $x_n = x_{n+2}x_{n+1}x_{n+2}^{-1}$
- (2) $x_{n+2} = x_1^{-1}x_2^{-1} \cdots x_{n-2}^{-1}x_{n-1}^{-1}x_{n+2}^{-1}x_{n+4}x_{n+2}x_{n+4}^{-1}x_{n+2}x_{n-1}x_{n-2} \cdots x_2x_1$
- (3) $(x_ix_{n+2})^2 = (x_{n+2}x_i)^2$, where $i = 1, \dots, n+1, n+4$
- (4) $[x_i, x_n] = e$, where $1 \leq i \leq n-1$
- (5) $[x_i, x_{n+2}x_nx_{n+2}^{-1}] = e$, where $1 \leq i \leq n-1$
- (6) $[x_i, x_{n+2}x_jx_{n+2}^{-1}] = e$, where $1 \leq i < j \leq n-1$
- (7) $[x_{n-1}^{-1}x_{n+1}x_{n-1}, x_i] = e$, where $1 \leq i \leq n-2$
- (8) $[x_{n+1}, x_{n+4}] = e$
- (9) $x_{n+4}(x_{n+2}x_n)^2x_{n-1} \cdots x_1 = e$.

We simplify now Relation (8) by substituting Relation (1) and Relation (9):

$$x_{n+2}^{-1}x_nx_{n+2} \cdot (x_{n+2}x_n)^2x_{n-1} \cdots x_1 = (x_{n+2}x_n)^2x_{n-1} \cdots x_1 \cdot x_{n+2}^{-1}x_nx_{n+2}.$$

Using Relations (3) for $i = n$, we get $[x_{n+2}x_nx_{n+2}^{-1}, x_{n-1} \cdots x_1] = e$, which is known already by Relations (5), and therefore is redundant. Moreover, using Relations (1), (3) and (4), Relations (7) (for $i = n-2$) can be translated to:

$$\begin{aligned} e &= [x_{n+1}, x_{n-1}x_{n-2}x_{n-1}^{-1}] = [x_{n+2}^{-1}x_nx_{n+2}, x_{n-1}x_{n-2}x_{n-1}^{-1}] = \\ &= [x_nx_{n+2}x_nx_{n+2}^{-1}x_n^{-1}, x_{n-1}x_{n-2}x_{n-1}^{-1}] = [x_{n+2}x_nx_{n+2}^{-1}, x_{n-1}x_{n-2}x_{n-1}^{-1}]. \end{aligned}$$

This relation is known already by Relations (5). If we check the other cases $i = n-3, \dots, 1$, we get $[x_{n+2}x_nx_{n+2}^{-1}, x_{n-1}x_ix_{n-1}^{-1}]$, for $1 \leq i \leq n-3$. These relations are known by Relations (5). Therefore, Relations (7) are redundant.

Notice that it is possible to combine Relations (5) and (6) to the general form $[x_i, x_{n+2}x_jx_{n+2}^{-1}] = e$, where $1 \leq i < j \leq n$. So we get:

- (1) $x_n = x_{n+2}x_{n+1}x_{n+2}^{-1}$
- (2) $x_{n+2} = x_1^{-1}x_2^{-1} \cdots x_{n-2}^{-1}x_{n-1}^{-1}x_{n+2}^{-1}x_{n+4}x_{n+2}x_{n+4}^{-1}x_{n+2}x_{n-1}x_{n-2} \cdots x_2x_1$
- (3) $(x_ix_{n+2})^2 = (x_{n+2}x_i)^2$, where $i = 1, \dots, n+1, n+4$
- (4) $[x_i, x_n] = e$, where $1 \leq i \leq n-1$
- (5) $[x_i, x_{n+2}x_jx_{n+2}^{-1}] = e$, where $1 \leq i < j \leq n$
- (6) $x_{n+4}(x_{n+2}x_n)^2x_{n-1} \cdots x_1 = e$.

Relation $(x_{n+1}x_{n+2})^2 = (x_{n+2}x_{n+1})^2$ (from Relations (3)) can be translated to $(x_nx_{n+2})^2 = (x_{n+2}x_n)^2$ by substituting Relation (1), and therefore is redundant.

Our goal now is to omit x_{n+4} . By Relations (3) for $i = n$ and Relations (4) and (6), Relation (3) for $i = n + 4$ can be rewritten as (by substituting x_{n+4} from Relation (2))

$$(x_{n+2} \cdot x_n x_{n-1} \cdots x_1)^2 = (x_n x_{n-1} \cdots x_1 \cdot x_{n+2})^2.$$

Using Relation (6) again, Relation (2) becomes:

$$\begin{aligned} x_{n+2} = & \quad x_1^{-1} \cdots x_{n-1}^{-1} x_{n+2}^{-1} (x_1^{-1} \cdots x_{n-1}^{-1} (x_{n+2} x_n)^{-2}) x_{n+2} \cdot \\ & \cdot ((x_{n+2} x_n)^2 x_{n-1} \cdots x_1) x_{n+2} x_{n-1} \cdots x_1, \end{aligned}$$

which can be written as $(x_{n+2} \cdot x_{n-1} \cdots x_1)^2 = (x_{n-1} \cdots x_1 \cdot x_{n+2})^2$. By Relation (1), the generator x_{n+1} is omitted. By Relation (6), x_{n+4} can be omitted too. Therefore, we have the following presentation:

Generators: $\{x_1, \dots, x_n, x_{n+2}\}$

Relations:

- (1) $(x_{n+2} \cdot x_{n-1} \cdots x_1)^2 = (x_{n-1} \cdots x_1 \cdot x_{n+2})^2$
- (2) $(x_i x_{n+2})^2 = (x_{n+2} x_i)^2$, where $1 \leq i \leq n$
- (3) $(x_{n+2} \cdot x_n x_{n-1} \cdots x_1)^2 = (x_n x_{n-1} \cdots x_1 \cdot x_{n+2})^2$
- (4) $[x_i, x_n] = e$, where $1 \leq i \leq n-1$
- (5) $[x_i, x_{n+2} x_j x_{n+2}^{-1}] = e$, where $1 \leq i < j \leq n$

Now, we show that Relations (1) and (3) are redundant. Let us denote $b := x_n$, $a := x_{n-1} \cdots x_1$ and $x := x_{n+2}$. In these notations, we have the relations $[a, b] = e$ (by Relations (4)), $(bx)^2 = (xb)^2$ (by Relations (2)), $(ax)^2 = (xa)^2$ (by Relation (1)) and $[xbx^{-1}, a] = e$ (by Relations (5)). We simplify Relation (3), which is $(bax)^2 = (xba)^2$. We do it as follows:

$$\begin{aligned} baxbax &= xba xba \rightarrow baxbx^{-1}xax = xba xbx^{-1}xa \rightarrow \\ &\rightarrow bxbx^{-1}axax = xbxbx^{-1}axa. \end{aligned}$$

The last resulting relation is redundant (that means that Relation (3) is redundant).

We show now that Relation (1) is redundant too. First, we write this relation in detail:

$$\begin{aligned}
 & x_{n+2} \cdot x_{n-1}x_{n-2} \cdots x_2x_1 \cdot x_{n+2} \cdot x_{n-1}x_{n-2} \cdots x_2x_1 = \\
 & = x_{n-1}x_{n-2} \cdots x_2x_1 \cdot x_{n+2} \cdot x_{n-1}x_{n-2} \cdots x_2x_1 \cdot x_{n+2}.
 \end{aligned}$$

We add the expression $x_{n+2}^{-1}x_{n+2} = e$:

$$\begin{aligned}
 & x_{n+2}x_{n-1}x_{n-2} \cdots x_2x_1(x_{n+2}x_{n-1}x_{n+2}^{-1})(x_{n+2}x_{n-2}x_{n+2}^{-1}) \cdots (x_{n+2}x_2x_{n+2}^{-1})x_{n+2}x_1 = \\
 & = x_{n-1}x_{n-2} \cdots x_2x_1(x_{n+2}x_{n-1}x_{n+2}^{-1})(x_{n+2}x_{n-2}x_{n+2}^{-1}) \cdots (x_{n+2}x_2x_{n+2}^{-1})x_{n+2}x_1x_{n+2}.
 \end{aligned}$$

By Relations (5), x_1 commutes with all expressions in the brackets. Therefore we get:

$$\begin{aligned}
 & x_{n+2}x_{n-1}x_{n-2} \cdots x_2(x_{n+2}x_{n-1}x_{n+2}^{-1})(x_{n+2}x_{n-2}x_{n+2}^{-1}) \cdots (x_{n+2}x_2x_{n+2}^{-1})x_1x_{n+2}x_1 = \\
 & = x_{n-1}x_{n-2} \cdots x_2(x_{n+2}x_{n-1}x_{n+2}^{-1})(x_{n+2}x_{n-2}x_{n+2}^{-1}) \cdots (x_{n+2}x_2x_{n+2}^{-1})x_1x_{n+2}x_1x_{n+2}.
 \end{aligned}$$

Since by Relations (2), $(x_1x_{n+2})^2 = (x_{n+2}x_1)^2$, we get:

$$\begin{aligned}
 & x_{n+2}x_{n-1}x_{n-2} \cdots x_2(x_{n+2}x_{n-1}x_{n+2}^{-1})(x_{n+2}x_{n-2}x_{n+2}^{-1}) \cdots (x_{n+2}x_3x_{n+2}^{-1})(x_{n+2}x_2x_{n+2}^{-1}) = \\
 & = x_{n-1}x_{n-2} \cdots x_2(x_{n+2}x_{n-1}x_{n+2}^{-1})(x_{n+2}x_{n-2}x_{n+2}^{-1}) \cdots (x_{n+2}x_3x_{n+2}^{-1})x_{n+2}x_2.
 \end{aligned}$$

Again by Relations (5), x_2 commutes with all the expressions in the brackets. Hence we have:

$$\begin{aligned}
 & x_{n+2}x_{n-1}x_{n-2} \cdots x_3(x_{n+2}x_{n-1}x_{n+2}^{-1})(x_{n+2}x_{n-2}x_{n+2}^{-1}) \cdots (x_{n+2}x_3x_{n+2}^{-1})x_2x_{n+2}x_2 = \\
 & = x_{n-1}x_{n-2} \cdots x_3(x_{n+2}x_{n-1}x_{n+2}^{-1})(x_{n+2}x_{n-2}x_{n+2}^{-1}) \cdots (x_{n+2}x_3x_{n+2}^{-1})x_2x_{n+2}x_2x_{n+2}.
 \end{aligned}$$

Since by Relations (2), $(x_2x_{n+2})^2 = (x_{n+2}x_2)^2$, we get a shorter form:

$$\begin{aligned}
 & x_{n+2}x_{n-1}x_{n-2} \cdots x_3(x_{n+2}x_{n-1}x_{n+2}^{-1})(x_{n+2}x_{n-2}x_{n+2}^{-1}) \cdots (x_{n+2}x_3x_{n+2}^{-1}) = \\
 & = x_{n-1}x_{n-2} \cdots x_3(x_{n+2}x_{n-1}x_{n+2}^{-1})(x_{n+2}x_{n-2}x_{n+2}^{-1}) \cdots x_{n+2}x_3.
 \end{aligned}$$

We continue in the same manner for $x_3, x_4, \dots, x_{n-2}$ in order to get the following relation:

$$x_{n+2}x_{n-1}(x_{n+2}x_{n-1}x_{n+2}^{-1})x_{n-2}x_{n+2}x_{n-2}x_{n+2}^{-1} = x_{n-1}(x_{n+2}x_{n-1}x_{n+2}^{-1})x_{n-2}x_{n+2}x_{n-2}.$$

Since $(x_{n-2}x_{n+2})^2 = (x_{n+2}x_{n-2})^2$ (by Relations (2)), we get $(x_{n-1}x_{n+2})^2 = (x_{n+2}x_{n-1})^2$, which appear already in Relations (2). Therefore, Relation (1) is redundant.

We finally get the requested presentation for $\pi_1(\mathbb{CP}^2 - T_{n,0})$ with $x_1, \dots, x_n, x_{n+2}$ as generators and with the following relations:

- (1) $(x_i x_{n+2})^2 = (x_{n+2} x_i)^2$, where $1 \leq i \leq n$
- (2) $[x_i, x_n] = e$, where $1 \leq i \leq n-1$
- (3) $[x_i, x_{n+2} x_j x_{n+2}^{-1}] = e$, where $1 \leq i < j \leq n$,

as needed. ■

By the above results, we can conclude:

Corollary 3.10. *The affine and projective fundamental groups of a conic-line arrangement, composed of two conics, which are tangent to each other at two points, and n tangent lines, are big.*

Proof. We have seen that the projective fundamental group of two tangent conics is $\langle a, b | (ab)^2 = (ba)^2 = e \rangle \cong \mathbb{Z} * \mathbb{Z}/2$, which is big, as we saw in Corollary 2.9.

Since the projective fundamental group is a quotient of the affine fundamental group by the projective relation, the affine fundamental group is big too.

Now, let A be a conic-line arrangement with two conics and with some additional tangent lines. Since the arrangement composed of two tangent conics is a sub-arrangement of A , and the fundamental groups of two tangent conics are big, we have that the fundamental groups of A are big too. ■

4. THE FUNDAMENTAL GROUP OF THE COMPLEMENT OF TWO TANGENT CONICS WITH $n + m$ TANGENT LINES

In this section, we compute the fundamental group of the complement of the arrangement $T_{n,m}$ consisting of two tangent conics, n lines

which are tangent to one of the conics and m lines which are tangent to the other one.

4.1. The braid monodromy factorizations. In this subsection we present the BMF of the arrangements $T_{1,1}$, $T_{2,1}$, $T_{2,2}$ and we shall compute the BMF of the arrangement $T_{n,m}$ for any n, m . Note that we dealt with the arrangement $T_{1,1}$ in [1], and we give here its BMF $\Delta_{T_{1,1}}^2$.

Lemma 4.1. *Let $T_{1,1}$ be a conic-line arrangement composed of two tangent conics (at two points) and two tangent lines, each is tangent to a different conic, see Figure 13.*





FIGURE 13. The arrangement $T_{1,1}$

Then its BMF is

$$\begin{aligned} \Delta_{T_{1,1}}^2 = & Z_{23}^2 \cdot \bar{Z}_{24}^2 \cdot (Z_{56})^{Z_{25}^{-2}} \cdot Z_{25}^4 \cdot Z_{46}^4 \cdot Z_{13}^4 \cdot (Z_{34})^{Z_{13}^2 Z_{46}^2} \cdot \\ & \cdot (Z_{16}^2)^{Z_{13}^2} \cdot (Z_{15}^2)^{Z_{12}^2} \cdot (Z_{34})^{Z_{45}^2} \cdot Z_{12}^2 \cdot Z_{45}^4 \cdot Z_{56}. \end{aligned}$$

In a similar way, one can compute the BMFs of $T_{2,1}$ and $T_{2,2}$:


Lemma 4.2. *Let $T_{2,1}$ be a conic-line arrangement composed of two tangent conics (at two points) and three additional lines, two of them are tangent to the first conic, and the third one is tangent to the second conic, see Figure 14.*

FIGURE 14. The arrangement $T_{2,1}$

Then its BMF is:

$$\begin{aligned} \Delta_{T_{2,1}}^2 = & Z_{23} \cdot Z_{67}^2 \cdot Z_{13}^4 \cdot Z_{24}^4 \cdot Z_{57}^4 \cdot (Z_{45})^{\bar{Z}_{24}^2 Z_{57}^2} \cdot (Z_{27}^2)^{Z_{24}^2 Z_{23}^2} \cdot (Z_{37}^2)^{Z_{13}^2} \cdot \\ & \cdot Z_{36}^4 \cdot (Z_{17}^2)^{Z_{13}^2} \cdot (Z_{45})^{Z_{34}^{-2} Z_{13}^2} \cdot (Z_{34}^4)^{Z_{13}^2} \cdot (Z_{15}^2)^{Z_{13}^2} \cdot (Z_{15}^2)^{Z_{13}^2} \cdot \\ & \cdot (Z_{23})^{Z_{36}^2 \bar{Z}_{13}^{-2}} \cdot Z_{46}^2 \cdot (Z_{56}^2)^{Z_{15}^2 Z_{13}^2} \cdot (Z_{16}^2)^{Z_{15}^2 Z_{13}^2}. \end{aligned}$$

Lemma 4.3. *Let $T_{2,2}$ be a conic-line arrangement composed of two tangent conics (at two points) and four additional lines, two of them are tangent to the first conic, and the two others are tangent to the second conic, see Figure 15.*

FIGURE 15. The arrangement $T_{2,2}$

Then its BMF is:

$$\begin{aligned}
 \Delta_{T_{2,2}}^2 = & Z_{23} \cdot Z_{67}^2 \cdot Z_{13}^4 \cdot Z_{24}^4 \cdot Z_{57}^4 \cdot (Z_{45})^{\bar{Z}_{24}^2 Z_{57}^2} \cdot (Z_{27}^2)^{Z_{24}^2 Z_{23}^2} \cdot (Z_{37}^2)^{Z_{13}^2} \cdot \\
 & \cdot (Z_{68}^2)^{Z_{67}^2} \cdot (\bar{Z}_{28}^2)^{\bar{Z}_{68}^2} \cdot (Z_{38}^2)^{Z_{37}^2 Z_{35}^2 Z_{13}^2} \cdot Z_{26}^4 \cdot (Z_{17}^2)^{Z_{13}^2} \cdot \\
 & \cdot (Z_{45})^{Z_{58}^{-2} Z_{78}^{-2} Z_{34}^{-2} Z_{13}^2} \cdot (Z_{58}^4)^{Z_{57}^2} \cdot (Z_{18}^2)^{Z_{17}^2 Z_{13}^2} \cdot Z_{78}^2 \cdot (Z_{15}^2)^{Z_{13}^2} \cdot \\
 & \cdot (Z_{34}^4)^{Z_{13}^2} \cdot (Z_{15}^2)^{Z_{13}^2} \cdot (Z_{23})^{Z_{12}^{-2} \bar{Z}_{26}^2} \cdot (Z_{46}^2)^{Z_{45}^2} \cdot (Z_{56}^2)^{Z_{15}^2 Z_{13}^2} \cdot (Z_{16}^2)^{Z_{15}^2 Z_{13}^2}.
 \end{aligned}$$


Now, we compute the BMF of $T_{n,m}$.

Proposition 4.4. *Let Q_1, Q_2 be two tangent conics in $\mathbb{CP}^2$. Consider two sets of n and m lines, such that n lines are tangent to Q_2 , and m lines are tangent to Q_1 . Denote: $T_{n,m} = Q_1 \cup Q_2 \cup (\bigcup_{i=1}^n L_i) \cup (\bigcup_{j=1}^m L'_j)$. Then:*

$$\begin{aligned}
 \Delta_{T_{n,m}}^2 = & Z_{23} \cdot (Z_{23})^{Z_{12}^{-2} \prod_{i=6}^{n+4} \bar{Z}_{2i}^2} \cdot (Z_{45})^{\bar{Z}_{24}^2 Z_{5,n+5}^2} \cdot \tilde{Z}_{45} \cdot \bar{Z}_{24}^4 \cdot (Z_{34}^4)^{Z_{13}^2} \cdot \\
 & \cdot Z_{13}^4 \cdot \prod_{i=6}^{n+4} \bar{Z}_{2i}^4 \cdot \prod_{i=n+5}^{n+m+4} (Z_{5i}^4)^{Z_{5,n+5}^2} \cdot (Z_{15}^2)^{Z_{13}^2} \cdot \prod_{i=6}^{n+4} \left[(Z_{4i}^2)^{Z_{45}^2} \cdot (Z_{5i}^2)^{Z_{15}^2 Z_{13}^2} \right] \cdot \\
 & \cdot \prod_{i=6}^{n+4} (Z_{1i}^2)^{Z_{15}^2 Z_{13}^2} \cdot \prod_{6 \leq i < j \leq n+4} (\bar{Z}_{ij}^2)^{\bar{Z}_{2i}^2} \cdot (Z_{2,n+5}^2)^{Z_{24}^2 Z_{23}^2} \cdot (Z_{3,n+5}^2)^{Z_{13}^2} \cdot \\
 & \cdot \prod_{i=n+6}^{n+m+4} \left[(\bar{Z}_{2i}^2)^{\left(\prod_{j=n+4}^6 \bar{Z}_{2j}^{-2} \right)} \cdot \tilde{Z}_{3i}^2 \right] \cdot \prod_{i=n+6}^{n+m+4} Z_{n+5,i}^2 \cdot \prod_{n+6 \leq i < j \leq n+m+4} (Z_{ij}^2)^{Z_{5i}^{-2} Z_{5,n+5}^2} \cdot \\
 & \cdot \prod_{i=n+5}^{n+m+4} (Z_{1i}^2)^{Z_{1,n+5}^2 Z_{13}^2} \cdot \prod_{i=6}^{n+4} \left[\prod_{j=n+5}^{n+m+4} (Z_{ij}^2)^{\left(\prod_{k=n+5}^j Z_{kj}^{-2} \right)} \right]
 \end{aligned}$$

The skeletons of $\tilde{Z}_{45}$ and $\tilde{Z}_{3i}^2$ appear in Figure 16.

Proof. By a similar technique to what we have used for computing the BMF of $T_{n,0}$, we compute the BMF of $T_{n,m}$.

FIGURE 16. Skeletons for the BMF of $T_{n,m}$

We first construct the BMF of $T_{1,m}$ by generalizing the BMF of $T_{1,1}$, $T_{1,2}$, $T_{1,3}$:

$$\begin{aligned}
\Delta_{T_{1,m}}^2 &= Z_{23} \cdot (Z_{23})^{Z_{12}^{-2}} \cdot (Z_{45})^{\bar{Z}_{24}^2 Z_{56}^2} \cdot \tilde{Z}'_{45} \cdot \bar{Z}_{24}^4 \cdot (Z_{34}^4)^{Z_{13}^2} \cdot Z_{13}^4 \cdot \prod_{i=6}^{m+5} (Z_{5i}^4)^{Z_{56}^2} \cdot \\
&\quad \cdot (Z_{15}^2)^{Z_{13}^2} \cdot (Z_{26}^2)^{Z_{24}^2 Z_{23}^2} \cdot (Z_{36}^2)^{Z_{13}^2} \cdot \left(\prod_{i=7}^{m+5} [\bar{Z}_{2i}^2 \cdot \tilde{Z}'_{3i}{}^2] \right) \cdot \prod_{i=7}^{m+5} Z_{6i}^2 \cdot \\
&\quad \cdot \prod_{7 \leq i < j \leq m+5} (Z_{ij}^2)^{Z_{5i}^{-2} Z_{56}^2} \cdot \prod_{i=6}^{m+5} (Z_{1i}^2)^{Z_{16}^2 Z_{13}^2}.
\end{aligned}$$

Then we proceed in the same manner to compute the BMF of $T_{2,m}$ and $T_{3,m}$, as presented in the following table. The table is constructed as follows: in each row we write the factors which are related to the

same type of singularities in the different arrangements.

	$\Delta_{T_{1,m}}^2$	$\Delta_{T_{2,m}}^2$	$\Delta_{T_{3,m}}^2$
(1)	Z_{23}	Z_{23}	Z_{23}
(2)	$(Z_{23})^{Z_{12}^{-2}}$	$(Z_{23})^{Z_{12}^{-2} \bar{Z}_{26}^2}$	$(Z_{23})^{Z_{12}^{-2} \bar{Z}_{26}^2 \bar{Z}_{27}^2}$
(3)	$(Z_{45})^{\bar{Z}_{24}^2 Z_{56}^2}$	$(Z_{45})^{\bar{Z}_{24}^2 Z_{57}^2}$	$(Z_{45})^{\bar{Z}_{24}^2 Z_{58}^2}$
(4)	$\tilde{Z}'_{45}$	$\tilde{Z}''_{45}$	$\tilde{Z}'''_{45}$
(5)	$\bar{Z}_{24}^4 \cdot (Z_{34})^{Z_{13}^2}$	$\bar{Z}_{24}^4 \cdot (Z_{34})^{Z_{13}^2}$	$\bar{Z}_{24}^4 \cdot (Z_{34})^{Z_{13}^2}$
(6)	Z_{13}^4	$Z_{13}^4 \cdot \bar{Z}_{26}^4$	$Z_{13}^4 \cdot \prod_{i=6,7} \bar{Z}_{2i}^4$
(7)	$\prod_{i=6}^{m+5} (Z_{5i}^4)^{Z_{56}^2}$	$\prod_{i=7}^{m+6} (Z_{5i}^4)^{Z_{57}^2}$	$\prod_{i=8}^{m+7} (Z_{5i}^4)^{Z_{58}^2}$
(8)	$(Z_{15}^2)^{Z_{13}^2}$	$(Z_{15}^2)^{Z_{13}^2} \cdot (Z_{46}^2)^{Z_{45}^2} \cdot (Z_{56}^2)^{Z_{15}^2 Z_{13}^2}$	$(Z_{15}^2)^{Z_{13}^2} \cdot \prod_{i=6}^7 [(Z_{4i}^2)^{Z_{45}^2} \cdot (Z_{5i}^2)^{Z_{15}^2 Z_{13}^2}]$
(9)	—	$(Z_{16}^2)^{Z_{15}^2 Z_{13}^2}$	$\prod_{i=6}^7 (Z_{1i}^2)^{Z_{15}^2 Z_{13}^2} \cdot (Z_{67}^2)^{\bar{Z}_{26}^2}$
(10)	$(Z_{26}^2)^{Z_{24}^2 Z_{23}^2} \cdot (Z_{36}^2)^{Z_{13}^2} \cdot \prod_{i=7}^{m+5} [\bar{Z}_{2i}^2 \cdot \tilde{Z}'_{3i}{}^2]$	$(Z_{27}^2)^{Z_{24}^2 Z_{23}^2} \cdot (Z_{37}^2)^{Z_{13}^2} \cdot \prod_{i=8}^{m+6} [\bar{Z}_{2i}^2 \cdot \tilde{Z}''_{3i}{}^2]$	$(Z_{28}^2)^{Z_{24}^2 Z_{23}^2} \cdot (Z_{38}^2)^{Z_{13}^2} \cdot \prod_{i=9}^{m+7} [\bar{Z}_{2i}^2 \cdot \tilde{Z}'''_{3i}{}^2]$
(11)	$\prod_{i=7}^{m+5} Z_{6i}^2 \cdot \prod_{7 \leq i < j \leq m+5} (Z_{ij}^2)^{Z_{5i}^{-2} Z_{56}^2}$	$\prod_{i=8}^{m+6} Z_{7i}^2 \cdot \prod_{8 \leq i < j \leq m+6} (Z_{ij}^2)^{Z_{5i}^{-2} Z_{57}^2}$	$\prod_{i=9}^{m+7} Z_{8i}^2 \cdot \prod_{9 \leq i < j \leq m+7} (Z_{ij}^2)^{Z_{5i}^{-2} Z_{58}^2}$
(12)	$\prod_{i=6}^{m+5} (Z_{1i}^2)^{Z_{16}^2 Z_{13}^2}$	$\prod_{i=7}^{m+6} (Z_{1i}^2)^{Z_{17}^2 Z_{13}^2} \cdot \prod_{j=7}^{m+6} (Z_{6j}^2)^{(\prod_{k=7}^j Z_{kj}^{-2})}$	$\prod_{i=8}^{m+7} (Z_{1i}^2)^{Z_{18}^2 Z_{13}^2} \cdot \prod_{i=6}^7 \left[\prod_{j=8}^{m+7} (Z_{ij}^2)^{(\prod_{k=8}^j Z_{kj}^{-2})} \right]$

The skeletons of $\tilde{Z}'_{45}$, $\tilde{Z}''_{45}$, $\tilde{Z}'''_{45}$ and $\tilde{Z}'_{3i}{}^2$, $\tilde{Z}''_{3i}{}^2$, $\tilde{Z}'''_{3i}{}^2$ appear in Figures 17 and 18 respectively.





FIGURE 17. The skeletons of $\tilde{Z}'_{45}$, $\tilde{Z}''_{45}$, $\tilde{Z}'''_{45}$

FIGURE 18. The skeletons of $\tilde{Z}'_{3i}{}^2$, $\tilde{Z}''_{3i}{}^2$, $\tilde{Z}'''_{3i}{}^2$

Rows (1) and (2) correspond to the branch points of Q_1 . Rows (3) and (4) correspond to the branch points of Q_2 . Row (5) corresponds to the two tangency points between Q_1 and Q_2 . Row (6) corresponds to the tangency points between Q_1 and the lines which are tangent to it. Row (7) corresponds to the tangency points between Q_2 and the lines which are tangent to it. Row (8) corresponds to the intersection points between the lines $\{L_i\}_{i=1}^n$ and the conic Q_2 . Row (9) corresponds to the intersection points between the lines $\{L_i\}_{i=1}^n$ themselves. Row (10) corresponds to the intersection points between the lines $\{L'_j\}_{j=1}^m$ with Q_1 . Row (11) corresponds to the intersection points between the lines $\{L'_j\}_{j=1}^m$ themselves. Row (12) corresponds to the intersection points between the lines $\{L_i\}_{i=1}^n$ and $\{L'_j\}_{j=1}^m$.

In the next step, we generalize the factorization for any n , i.e. we compute $\Delta_{T_{n,m}}^2$. In the following table, each row includes the general

form of the corresponding row in the previous table:

	$\Delta_{T_{n,m}}^2$
(1)	Z_{23}
(2)	$(Z_{23})^{Z_{12}^{-2} \prod_{i=6}^{n+4} \bar{Z}_{2i}^2}$
(3)	$(Z_{45})^{Z_{24}^2 Z_{5,n+5}^2}$
(4)	$\tilde{Z}_{45}$
(5)	$\bar{Z}_{24}^4 \cdot (Z_{34}^4)^{Z_{13}^2}$
(6)	$Z_{13}^4 \cdot \prod_{i=6}^{n+4} \bar{Z}_{2i}^4$
(7)	$\prod_{i=n+5}^{n+m+4} (Z_{5i}^4)^{Z_{5,n+5}^2}$
(8)	$(Z_{15}^2)^{Z_{13}^2} \cdot \prod_{i=6}^{n+4} \left[(Z_{4i}^2)^{Z_{45}^2} \cdot (Z_{5i}^2)^{Z_{15}^2 Z_{13}^2} \right]$
(9)	$\prod_{i=6}^{n+4} (Z_{1i}^2)^{Z_{15}^2 Z_{13}^2} \cdot \prod_{6 \leq i < j \leq n+4} (\bar{Z}_{ij}^2)^{\bar{Z}_{2i}^2}$
(10)	$(Z_{2,n+5}^2)^{Z_{24}^2 Z_{23}^2} \cdot (Z_{3,n+5}^2)^{Z_{13}^2} \cdot \prod_{i=n+6}^{n+m+4} \left[(\bar{Z}_{2i}^2)^{\left(\prod_{j=n+4}^6 \bar{Z}_{2j}^{-2} \right)} \cdot \tilde{Z}_{3i}^2 \right]$
(11)	$\prod_{i=n+6}^{n+m+4} Z_{n+5,i}^2 \cdot \prod_{n+6 \leq i < j \leq n+m+4} (Z_{ij}^2)^{Z_{5i}^{-2} Z_{5,n+5}^2}$
(12)	$\prod_{i=n+5}^{n+m+4} (Z_{1i}^2)^{Z_{1,n+5}^2 Z_{13}^2} \cdot \prod_{i=6}^{n+4} \left[\prod_{j=n+5}^{n+m+4} (Z_{ij}^2)^{\left(\prod_{k=n+5}^j Z_{kj}^{-2} \right)} \right]$

The skeletons of $\tilde{Z}_{45}$ and $\tilde{Z}_{3i}^2$ appear in Figure 16. ■

4.2. The corresponding fundamental groups. In this section, we compute the fundamental group of the complement of the general arrangement $T_{n,m}$, $n, m \geq 1$.

Note that the arrangement $T_{1,1}$ appears already in [1], therefore we cite here the presentation of its fundamental group:

Proposition 4.5. *Let $T_{1,1}$ be the curve defined in Lemma 4.1, see Figure 13.*

Then:

$$\pi_1(\mathbb{CP}^2 - T_{1,1}) \cong \langle x_1 \rangle \oplus \langle x_2, x_3 \mid (x_2 x_3)^2 = (x_3 x_2)^2 \rangle.$$

Now, we prove Theorem 1.1 by computing the simplified presentation of the fundamental group of $T_{n,m}$:

Proof of Theorem 1.1:

Using the van Kampen Theorem on $\Delta_{T_{n,m}}^2$ (computed in Proposition 4.4), we get the following presentation for $\pi_1(\mathbb{CP}^2 - T_{n,m})$:

Generators: $\{x_1, x_2, x_3, x_4, x_5, x_6, x_7, \dots, x_{n+4}, x_{n+5}, \dots, x_{n+m+4}\}$.

Relations:

- (1) $x_2 = x_3$
- (2) $x_3 = x_1^{-1} x_3^{-1} x_4^{-1} x_5^{-1} \cdot x_{n+4} x_{n+3} \cdots x_3 x_2 x_3^{-1} \cdots x_{n+3}^{-1} x_{n+4}^{-1} \cdot x_5 x_4 x_3 x_1$
- (3) $x_4 x_3 x_2 x_3^{-1} x_4 x_3 x_2^{-1} x_3^{-1} x_4^{-1} = x_{n+5} x_5 x_{n+5}^{-1}$
- (4) $x_3 x_1 x_3^{-1} x_1^{-1} x_3^{-1} x_4 x_3 x_1 x_3 x_1^{-1} x_3^{-1} = x_5^{-1} \cdot x_{n+5}^{-1} x_{n+6}^{-1} \cdots x_{n+m+4}^{-1} \cdot x_{n+5} x_5 x_{n+5}^{-1} \cdot x_{n+m+4} \cdots x_{n+6} x_{n+5} \cdot x_5$
- (5) $(x_3 x_2 x_3^{-1} x_4)^2 = (x_4 x_3 x_2 x_3^{-1})^2$
- (6) $(x_3 x_1 x_3 x_1^{-1} x_3^{-1} x_4)^2 = (x_4 x_3 x_1 x_3 x_1^{-1} x_3^{-1})^2$
- (7) $(x_1 x_3)^2 = (x_3 x_1)^2$
- (8) $(x_2 \cdot x_3^{-1} x_4^{-1} \cdots x_{i-1}^{-1} x_i x_{i-1} \cdots x_4 x_3)^2 = (x_3^{-1} x_4^{-1} \cdots x_{i-1}^{-1} x_i x_{i-1} \cdots x_4 x_3 \cdot x_2)^2$,
where $6 \leq i \leq n+4$
- (9) $(x_5 x_{n+5})^2 = (x_{n+5} x_5)^2$
- (10) $(x_5 x_{n+5}^{-1} x_i x_{n+5})^2 = (x_{n+5}^{-1} x_i x_{n+5} x_5)^2$, where $n+6 \leq i \leq n+m+4$
- (11) $[x_3 x_1 x_3^{-1}, x_5] = e$
- (12) $[x_5 x_4 x_5^{-1}, x_i] = e$, where $6 \leq i \leq n+4$
- (13) $[x_5 x_3 x_1 x_3^{-1} x_5 x_3 x_1^{-1} x_3^{-1} x_5^{-1}, x_i] = e$, where $6 \leq i \leq n+4$
- (14) $[x_5 x_3 x_1 x_3^{-1} x_5^{-1}, x_i] = e$, where $6 \leq i \leq n+4$.
- (15) $[x_{j-1} x_{j-2} \cdots x_4 x_3 x_2 x_3^{-1} x_4^{-1} \cdots x_{i-2}^{-1} x_{i-1}^{-1} x_i x_{i-1} x_{i-2} \cdots x_4 x_3 x_2^{-1} x_3^{-1} x_4^{-1} \cdots x_{j-2}^{-1} x_{j-1}^{-1}, x_j] = e$, where $6 \leq i < j \leq n+4$
- (16) $[x_4 x_3 x_2 x_3^{-1} x_4^{-1}, x_{n+5}] = e$
- (17) $[x_3 x_1 x_3 x_1^{-1} x_3^{-1}, x_{n+5}] = e$
- (18) $[x_5 x_4 x_3 x_2 x_3^{-1} x_4^{-1} x_5^{-1}, x_{n+5} x_{n+6}^{-1} \cdots x_{i-2}^{-1} x_{i-1}^{-1} x_i x_{i-1} x_{i-2} \cdots x_{n+6} x_{n+5}] = e$,
where $n+6 \leq i \leq n+m+4$
- (19) $[x_5 x_3 x_1 x_3 x_1^{-1} x_3^{-1} x_5^{-1}, x_{n+5} x_{n+6}^{-1} \cdots x_{i-2}^{-1} x_{i-1}^{-1} x_i x_{i-1} x_{i-2} \cdots x_{n+6} x_{n+5}] = e$,
where $n+6 \leq i \leq n+m+4$
- (20) $[x_{n+5}, x_i] = e$, where $n+6 \leq i \leq n+m+4$
- (21) $[x_{n+5} x_5^{-1} x_{n+5}^{-1} x_i x_{n+5} x_5 x_{n+5}^{-1}, x_j] = e$, where $n+6 \leq i < j \leq n+m+4$
- (22) $[x_3 x_1 x_3^{-1}, x_{n+5}^{-1} x_i x_{n+5}] = e$, where $n+5 \leq i \leq n+m+4$
- (23) $[x_i, x_{n+5} x_{n+6}^{-1} \cdots x_{j-1}^{-1} x_j x_{j-1} \cdots x_{n+6} x_{n+5}] = e$, where $6 \leq i \leq n+4$ and $n+5 \leq j \leq n+m+4$
- (24) $x_{n+m+4} x_{n+m+3} \cdots x_4 x_3 x_2 x_1 = e$. (Projective relation).

Since $x_2 = x_3$ and $(x_1 x_3)^2 = (x_3 x_1)^2$ (by Relations (1) and (7)), we derive a simpler presentation as follows (we combine Relations (5) and (7) into one set):

Generators: $\{x_1, x_2, x_4, x_5, x_6, x_7, \dots, x_{n+4}, x_{n+5}, \dots, x_{n+m+4}\}$.

Relations:

- (1) $x_1^{-1} x_2 x_1 = x_4^{-1} x_5^{-1} \cdot x_{n+4} x_{n+3} \cdots x_4 x_2 x_4^{-1} \cdots x_{n+3}^{-1} x_{n+4}^{-1} \cdot x_5 x_4$

- (2) $x_4x_2x_4x_2^{-1}x_4^{-1} = x_{n+5}x_5x_{n+5}^{-1}$
- (3) $x_1^{-1}x_2^{-1}x_1x_4x_1^{-1}x_2x_1 = x_5^{-1} \cdot x_{n+5}^{-1}x_{n+6}^{-1} \cdots x_{n+m+4}^{-1} \cdot x_{n+5}x_5x_{n+5}^{-1} \cdot x_{n+m+4} \cdots x_{n+6}x_{n+5} \cdot x_5$
- (4) $(x_2x_i)^2 = (x_ix_2)^2$, where $i = 1, 4$
- (5) $(x_1^{-1}x_2x_1x_4)^2 = (x_4x_1^{-1}x_2x_1)^2$
- (6) $(x_4^{-1} \cdots x_{i-1}^{-1}x_ix_{i-1} \cdots x_4x_2)^2 = (x_2^{-1}x_4^{-1} \cdots x_{i-1}^{-1}x_ix_{i-1} \cdots x_4x_2^2)^2$,
where $6 \leq i \leq n+4$
- (7) $(x_5x_{n+5})^2 = (x_{n+5}x_5)^2$
- (8) $(x_5x_{n+5}^{-1}x_ix_{n+5})^2 = (x_{n+5}^{-1}x_ix_{n+5}x_5)^2$, where $n+6 \leq i \leq n+m+4$
- (9) $[x_2x_1x_2^{-1}, x_5] = e$
- (10) $[x_5x_4x_5^{-1}, x_i] = e$, where $6 \leq i \leq n+4$
- (11) $[x_5x_2x_1x_2^{-1}x_5x_2x_1^{-1}x_2^{-1}x_5^{-1}, x_i] = e$, where $6 \leq i \leq n+4$
- (12) $[x_5x_2x_1x_2^{-1}x_5^{-1}, x_i] = e$, where $6 \leq i \leq n+4$.
- (13) $[x_{j-1}x_{j-2} \cdots x_4x_2x_4^{-1} \cdots x_{i-2}^{-1}x_{i-1}^{-1}x_ix_{i-1}x_{i-2} \cdots x_4x_2^{-1}x_4^{-1} \cdots x_{j-2}^{-1}x_{j-1}^{-1}, x_j] = e$,
where $6 \leq i < j \leq n+4$
- (14) $[x_4x_2x_4^{-1}, x_{n+5}] = e$
- (15) $[x_1^{-1}x_2x_1, x_{n+5}] = e$
- (16) $[x_5x_4x_2x_4^{-1}x_5^{-1}, x_{n+5}x_{n+6} \cdots x_{i-2}^{-1}x_{i-1}^{-1}x_ix_{i-1}x_{i-2} \cdots x_{n+6}x_{n+5}] = e$,
where $n+6 \leq i \leq n+m+4$
- (17) $[x_5x_1^{-1}x_2x_1x_5^{-1}, x_{n+5}x_{n+6} \cdots x_{i-2}^{-1}x_{i-1}^{-1}x_ix_{i-1}x_{i-2} \cdots x_{n+6}x_{n+5}] = e$,
where $n+6 \leq i \leq n+m+4$
- (18) $[x_{n+5}, x_i] = e$, where $n+6 \leq i \leq n+m+4$
- (19) $[x_{n+5}x_5^{-1}x_{n+5}^{-1}x_ix_{n+5}x_5x_{n+5}^{-1}, x_j] = e$, where $n+6 \leq i < j \leq n+m+4$
- (20) $[x_2x_1x_2^{-1}, x_{n+5}x_ix_{n+5}] = e$, where $n+5 \leq i \leq n+m+4$
- (21) $[x_i, x_{n+5}x_{n+6} \cdots x_{j-1}^{-1}x_jx_{j-1} \cdots x_{n+6}x_{n+5}] = e$,
where $6 \leq i \leq n+4$ and $n+5 \leq j \leq n+m+4$
- (22) $x_{n+m+4}x_{n+m+3} \cdots x_5x_4x_2^2x_1 = e$.

Now we apply Relations (4) and (18) to get a much convenient presentation (we also combine Relations (7) and (8) into one set):

- (1) $x_1^{-1}x_2x_1 = x_4^{-1}x_5^{-1} \cdot x_{n+4}x_{n+3} \cdots x_4x_2x_4^{-1} \cdots x_{n+3}^{-1}x_{n+4}^{-1} \cdot x_5x_4$
- (2) $x_2^{-1}x_4x_2 = x_{n+5}x_5x_{n+5}^{-1}$
- (3) $x_1^{-1}x_2^{-1}x_1x_4x_1^{-1}x_2x_1 = x_5^{-1} \cdot x_{n+6}^{-1} \cdots x_{n+m+4}^{-1} \cdot x_5 \cdot x_{n+m+4} \cdots x_{n+6} \cdot x_5$
- (4) $(x_2x_i)^2 = (x_ix_2)^2$, where $i = 1, 4$
- (5) $(x_1^{-1}x_2x_1x_4)^2 = (x_4x_1^{-1}x_2x_1)^2$
- (6) $(x_4^{-1} \cdots x_{i-1}^{-1}x_ix_{i-1} \cdots x_4x_2)^2 = (x_2^{-1}x_4^{-1} \cdots x_{i-1}^{-1}x_ix_{i-1} \cdots x_4x_2^2)^2$,
where $6 \leq i \leq n+4$
- (7) $(x_5x_i)^2 = (x_ix_5)^2$, where $n+5 \leq i \leq n+m+4$
- (8) $[x_2x_1x_2^{-1}, x_5] = e$
- (9) $[x_5x_4x_5^{-1}, x_i] = e$, where $6 \leq i \leq n+4$
- (10) $[x_5x_2x_1x_2^{-1}x_5x_2x_1^{-1}x_2^{-1}x_5^{-1}, x_i] = e$, where $6 \leq i \leq n+4$
- (11) $[x_5x_2x_1x_2^{-1}x_5^{-1}, x_i] = e$, where $6 \leq i \leq n+4$.
- (12) $[x_{j-1}x_{j-2} \cdots x_4x_2x_4^{-1} \cdots x_{i-2}^{-1}x_{i-1}^{-1}x_ix_{i-1}x_{i-2} \cdots x_4x_2^{-1}x_4^{-1} \cdots x_{j-2}^{-1}x_{j-1}^{-1}, x_j] = e$,
where $6 \leq i < j \leq n+4$
- (13) $[x_4x_2x_4^{-1}, x_{n+5}] = e$
- (14) $[x_1^{-1}x_2x_1, x_{n+5}] = e$
- (15) $[x_5x_4x_2x_4^{-1}x_5^{-1}, x_{n+6} \cdots x_{i-2}^{-1}x_{i-1}^{-1}x_ix_{i-1}x_{i-2} \cdots x_{n+6}] = e$, where $n+6 \leq i \leq n+m+4$
- (16) $[x_5x_1^{-1}x_2x_1x_5^{-1}, x_{n+6} \cdots x_{i-2}^{-1}x_{i-1}^{-1}x_ix_{i-1}x_{i-2} \cdots x_{n+6}] = e$, where $n+6 \leq i \leq n+m+4$

- (17) $[x_{n+5}, x_i] = e$, where $n+6 \leq i \leq n+m+4$
- (18) $[x_5^{-1}x_ix_5, x_j] = e$, where $n+6 \leq i < j \leq n+m+4$
- (19) $[x_2x_1x_2^{-1}, x_i] = e$, where $n+5 \leq i \leq n+m+4$
- (20) $[x_i, x_{n+6}^{-1} \cdots x_{j-1}^{-1}x_jx_{j-1} \cdots x_{n+6}] = e$, where $6 \leq i \leq n+4$ and $n+5 \leq j \leq n+m+4$
- (21) $x_{n+m+4}x_{n+m+3} \cdots x_5x_4x_2^2x_1 = e$.

By Relations (9) we get the following simplified relations (we combine Relations (8), (11), (19) into one set):

- (1) $x_1^{-1}x_2x_1 = x_4^{-1}x_5^{-1} \cdot x_{n+4}x_{n+3} \cdots x_4x_2x_4^{-1} \cdots x_{n+3}^{-1}x_{n+4}^{-1} \cdot x_5x_4$
- (2) $x_2^{-1}x_4x_2 = x_{n+5}x_5x_{n+5}^{-1}$
- (3) $x_1^{-1}x_2^{-1}x_1x_4x_1^{-1}x_2x_1 = x_5^{-1} \cdot x_{n+6}^{-1} \cdots x_{n+m+4}^{-1} \cdot x_5 \cdot x_{n+m+4} \cdots x_{n+6} \cdot x_5$
- (4) $(x_2x_i)^2 = (x_ix_2)^2$, where $i = 1, 4$
- (5) $(x_1^{-1}x_2x_1x_4)^2 = (x_4x_1^{-1}x_2x_1)^2$
- (6) $(x_4^{-1} \cdots x_{i-1}^{-1}x_ix_{i-1} \cdots x_4x_2)^2 = (x_2^{-1}x_4^{-1} \cdots x_{i-1}^{-1}x_ix_{i-1} \cdots x_4x_2^2)^2$,
where $6 \leq i \leq n+4$
- (7) $(x_5x_i)^2 = (x_ix_5)^2$, where $n+5 \leq i \leq n+m+4$
- (8) $[x_2x_1x_2^{-1}, x_i] = e$, where $5 \leq i \leq n+m+4$
- (9) $[x_5x_4x_5^{-1}, x_i] = e$, where $6 \leq i \leq n+4$
- (10) $[x_5, x_i] = e$, where $6 \leq i \leq n+4$
- (11) $[x_{j-1}x_{j-2} \cdots x_4x_2x_4^{-1} \cdots x_{i-2}^{-1}x_{i-1}^{-1}x_ix_{i-1}x_{i-2} \cdots x_4x_2^{-1}x_4^{-1} \cdots x_{j-2}^{-1}x_{j-1}^{-1}, x_j] = e$,
where $6 \leq i < j \leq n+4$
- (12) $[x_4x_2x_4^{-1}, x_{n+5}] = e$
- (13) $[x_1^{-1}x_2x_1, x_{n+5}] = e$
- (14) $[x_5x_4x_2x_4^{-1}x_5^{-1}, x_{n+6}^{-1} \cdots x_{i-2}^{-1}x_{i-1}^{-1}x_ix_{i-1}x_{i-2} \cdots x_{n+6}] = e$, where $n+6 \leq i \leq n+m+4$
- (15) $[x_5x_1^{-1}x_2x_1x_5^{-1}, x_{n+6}^{-1} \cdots x_{i-2}^{-1}x_{i-1}^{-1}x_ix_{i-1}x_{i-2} \cdots x_{n+6}] = e$, where $n+6 \leq i \leq n+m+4$
- (16) $[x_{n+5}, x_i] = e$, where $n+6 \leq i \leq n+m+4$
- (17) $[x_5^{-1}x_ix_5, x_j] = e$, where $n+6 \leq i < j \leq n+m+4$
- (18) $[x_i, x_{n+6}^{-1} \cdots x_{j-1}^{-1}x_jx_{j-1} \cdots x_{n+6}] = e$, where $6 \leq i \leq n+4$ and $n+5 \leq j \leq n+m+4$
- (19) $x_{n+m+4}x_{n+m+3} \cdots x_5x_4x_2^2x_1 = e$.

By Relations (10), Relations (9) become $[x_4, x_i] = e$, where $6 \leq i \leq n+4$. These relations enable us to proceed in simplification:

- (1) $x_1^{-1}x_2x_1 = x_{n+4}x_{n+3} \cdots x_6x_2x_6^{-1} \cdots x_{n+3}^{-1}x_{n+4}^{-1}$
- (2) $x_2^{-1}x_4x_2 = x_{n+5}x_5x_{n+5}^{-1}$
- (3) $x_1^{-1}x_2^{-1}x_1x_4x_1^{-1}x_2x_1 = x_5^{-1} \cdot x_{n+6}^{-1} \cdots x_{n+m+4}^{-1} \cdot x_5 \cdot x_{n+m+4} \cdots x_{n+6} \cdot x_5$
- (4) $(x_2x_i)^2 = (x_ix_2)^2$, where $i = 1, 4$
- (5) $(x_1^{-1}x_2x_1x_4)^2 = (x_4x_1^{-1}x_2x_1)^2$
- (6) $(x_6^{-1} \cdots x_{i-1}^{-1}x_ix_{i-1} \cdots x_6x_2)^2 = (x_2^{-1}x_6^{-1} \cdots x_{i-1}^{-1}x_ix_{i-1} \cdots x_6x_2^2)^2$,
where $6 \leq i \leq n+4$
- (7) $(x_5x_i)^2 = (x_ix_5)^2$, where $n+5 \leq i \leq n+m+4$
- (8) $[x_2x_1x_2^{-1}, x_i] = e$, where $5 \leq i \leq n+m+4$
- (9) $[x_4, x_i] = e$, where $6 \leq i \leq n+4$
- (10) $[x_5, x_i] = e$, where $6 \leq i \leq n+4$
- (11) $[x_{j-1}x_{j-2} \cdots x_6x_2x_6^{-1} \cdots x_{i-2}^{-1}x_{i-1}^{-1}x_ix_{i-1}x_{i-2} \cdots x_6x_2^{-1}x_6^{-1} \cdots x_{j-2}^{-1}x_{j-1}^{-1}, x_j] = e$,
where $6 \leq i < j \leq n+4$
- (12) $[x_4x_2x_4^{-1}, x_{n+5}] = e$
- (13) $[x_1^{-1}x_2x_1, x_{n+5}] = e$
- (14) $[x_5x_4x_2x_4^{-1}x_5^{-1}, x_{n+6}^{-1} \cdots x_{i-2}^{-1}x_{i-1}^{-1}x_ix_{i-1}x_{i-2} \cdots x_{n+6}] = e$, where $n+6 \leq i \leq n+m+4$

- (15) $[x_5x_1^{-1}x_2x_1x_5^{-1}, x_{n+6}^{-1} \cdots x_{i-2}^{-1}x_{i-1}^{-1}x_ix_{i-1}x_{i-2} \cdots x_{n+6}] = e$, where $n+6 \leq i \leq n+m+4$
- (16) $[x_{n+5}, x_i] = e$, where $n+6 \leq i \leq n+m+4$
- (17) $[x_5^{-1}x_ix_5, x_j] = e$, where $n+6 \leq i < j \leq n+m+4$
- (18) $[x_i, x_{n+6}^{-1} \cdots x_{j-1}^{-1}x_jx_{j-1} \cdots x_{n+6}] = e$, where $6 \leq i \leq n+4$ and $n+5 \leq j \leq n+m+4$
- (19) $x_{n+m+4}x_{n+m+3} \cdots x_5x_4x_2^2x_1 = e$.

We treat Relations (18). Substituting $i = 6$ and $j = n+5$, we get $[x_6, x_{n+5}] = e$. Substituting $i = 6$ and $j = n+6$, we get $[x_6, x_{n+6}] = e$. If we take $i = 6$ and $j = n+7$, we get $[x_6, x_{n+6}^{-1}x_{n+7}x_{n+6}] = e$, which is $[x_6, x_{n+7}] = e$. Continuing this process by substituting $i = 6$ and $n+8 \leq j \leq n+m+4$, we get $[x_6, x_j] = e$ for $n+8 \leq j \leq n+m+4$. Now we substitute $i = 7$ and $n+5 \leq j \leq n+m+4$. From these substitutions we get: $[x_7, x_j] = e$ for $n+5 \leq j \leq n+m+4$. In the same manner, we get $[x_i, x_j] = e$ for $8 \leq i \leq n+4$ and $n+5 \leq j \leq n+m+4$. This enables us to simplify the above list of relations (we combine Relations (9), (10) and (18) into one set):

- (1) $x_1^{-1}x_2x_1 = x_{n+4}x_{n+3} \cdots x_6x_2x_6^{-1} \cdots x_{n+3}^{-1}x_{n+4}^{-1}$
- (2) $x_2^{-1}x_4x_2 = x_{n+5}x_5x_{n+5}^{-1}$
- (3) $x_1^{-1}x_2^{-1}x_1x_4x_1^{-1}x_2x_1 = x_5^{-1} \cdot x_{n+6}^{-1} \cdots x_{n+m+4}^{-1} \cdot x_5 \cdot x_{n+m+4} \cdots x_{n+6} \cdot x_5$
- (4) $(x_2x_i)^2 = (x_ix_2)^2$, where $i = 1, 4$
- (5) $(x_1^{-1}x_2x_1x_4)^2 = (x_4x_1^{-1}x_2x_1)^2$
- (6) $(x_6^{-1} \cdots x_{i-1}^{-1}x_ix_{i-1} \cdots x_6x_2)^2 = (x_2^{-1}x_6^{-1} \cdots x_{i-1}^{-1}x_ix_{i-1} \cdots x_6x_2^2)^2$,
where $6 \leq i \leq n+4$
- (7) $(x_5x_i)^2 = (x_ix_5)^2$, where $n+5 \leq i \leq n+m+4$
- (8) $[x_2x_1x_2^{-1}, x_i] = e$, where $5 \leq i \leq n+m+4$
- (9) $[x_i, x_j] = e$, where $6 \leq i \leq n+4$ and $j = 4, 5, n+5, \dots, n+m+4$
- (10) $[x_{j-1}x_{j-2} \cdots x_6x_2x_6^{-1} \cdots x_{i-2}^{-1}x_{i-1}^{-1}x_ix_{i-1}x_{i-2} \cdots x_6x_2^{-1}x_6^{-1} \cdots x_{j-2}^{-1}x_{j-1}^{-1}, x_j] = e$,
where $6 \leq i < j \leq n+4$
- (11) $[x_4x_2x_4^{-1}, x_{n+5}] = e$
- (12) $[x_1^{-1}x_2x_1, x_{n+5}] = e$
- (13) $[x_5x_4x_2x_4^{-1}x_5^{-1}, x_{n+6}^{-1} \cdots x_{i-2}^{-1}x_{i-1}^{-1}x_ix_{i-1}x_{i-2} \cdots x_{n+6}] = e$, where $n+6 \leq i \leq n+m+4$
- (14) $[x_5x_1^{-1}x_2x_1x_5^{-1}, x_{n+6}^{-1} \cdots x_{i-2}^{-1}x_{i-1}^{-1}x_ix_{i-1}x_{i-2} \cdots x_{n+6}] = e$, where $n+6 \leq i \leq n+m+4$
- (15) $[x_{n+5}, x_i] = e$, where $n+6 \leq i \leq n+m+4$
- (16) $[x_5^{-1}x_ix_5, x_j] = e$, where $n+6 \leq i < j \leq n+m+4$
- (17) $x_{n+m+4}x_{n+m+3} \cdots x_5x_4x_2^2x_1 = e$.

Now we simplify Relations (6) and (10) (we have used this trick of simplification in the computation of $\pi_1(\mathbb{C}^2 - C_n)$ in Section 2).

We start by substituting $i = 6$ in Relations (6) to get $(x_2x_6)^2 = (x_6x_2)^2$. We continue with $i = 6, j = 7$ in Relations (10) to get $[x_2^{-1}x_6x_2, x_7] = e$ (using $(x_2x_6)^2 = (x_6x_2)^2$). Using these two relations, we simplify Relations (6) for $i = 7$: $(x_2x_6^{-1}x_7x_6)^2 = (x_6^{-1}x_7x_6x_2)^2$. We

add the expression $x_2^{-1}x_2 = e$ in four locations as follows:

$$x_6x_2x_6^{-1}(x_2^{-1}x_2)x_7x_6x_2x_6^{-1}(x_2^{-1}x_2)x_7 = x_7x_6x_2x_6^{-1}(x_2^{-1}x_2)x_7x_6x_2x_6^{-1}(x_2^{-1}x_2).$$

Since $(x_2x_6)^2 = (x_6x_2)^2$, we can rewrite the relation as:

$$x_2^{-1}x_6^{-1}x_2x_6x_2x_7x_2^{-1}x_6^{-1}x_2x_6x_2x_7 = x_7x_2^{-1}x_6^{-1}x_2x_6x_2x_7x_2^{-1}x_6^{-1}x_2x_6x_2.$$

We use the relation $[x_2^{-1}x_6x_2, x_7] = e$ to get:

$$x_2^{-1}x_6^{-1}x_2x_6x_2x_2^{-1}x_6^{-1}x_2x_7x_6x_2x_7 = x_2^{-1}x_6^{-1}x_2x_7x_6x_2x_2^{-1}x_6^{-1}x_2x_7x_6x_2,$$

namely $x_2x_7x_6x_2x_7 = x_7x_2x_7x_6x_2$. We add the expression $x_2x_2^{-1} = e$ to get $x_2x_7(x_2x_2^{-1})x_6x_2x_7 = x_7x_2x_7x_6x_2$, and using $[x_2^{-1}x_6x_2, x_7] = e$, we get $(x_2x_7)^2 = (x_7x_2)^2$.

Now we proceed with $i = 6, j = 8$ and then $i = 7, j = 8$ in Relations (10) to get $[x_2^{-1}x_6x_2, x_8] = e$ and $[x_2^{-1}x_7x_2, x_8] = e$ respectively. These relations enable us to simplify the relation $(x_2x_6^{-1}x_7^{-1}x_8x_7x_6)^2 = (x_6^{-1}x_7^{-1}x_8x_7x_6x_2)^2$ (which we get from Relations (6) for $i = 8$) to $(x_2x_8)^2 = (x_8x_2)^2$.

In the same manner, we conclude that Relations (6) and (10) can be simplified to $(x_2x_i)^2 = (x_ix_2)^2$ for $6 \leq i \leq n+4$ and $[x_2^{-1}x_ix_2, x_j] = e$ for $6 \leq i < j \leq n+4$ respectively.

Relation (12) gets the form $[x_2, x_{n+5}] = e$ by substituting Relation (1) in it and by using Relations (9).

Relations (14) can be rewritten by substituting Relation (1) in them (for $n+6 \leq i \leq n+m+4$):

$$[x_5x_{n+4}x_{n+3} \cdots x_6x_2x_6^{-1} \cdots x_{n+3}^{-1}x_{n+4}^{-1}x_5^{-1}, x_{n+6}^{-1} \cdots x_{i-2}^{-1}x_{i-1}^{-1}x_ix_{i-1}x_{i-2} \cdots x_{n+6}] = e.$$

Now, by Relations (9), we get:

$$[x_5x_2x_5^{-1}, x_{n+6}^{-1} \cdots x_{i-2}^{-1}x_{i-1}^{-1}x_ix_{i-1}x_{i-2} \cdots x_{n+6}] = e.$$

For $i = n+6$, we get $[x_5x_2x_5^{-1}, x_{n+6}] = e$. Now, for $i = n+7$, we get $[x_5x_2x_5^{-1}, x_{n+6}^{-1}x_{n+7}x_{n+6}] = e$. By $[x_5x_2x_5^{-1}, x_{n+6}] = e$, it is simplified to $[x_5x_2x_5^{-1}, x_{n+7}] = e$. In the same way, we get that Relations (14) are equivalent to $[x_5x_2x_5^{-1}, x_i] = e$, for $n+6 \leq i \leq n+m+4$.

Therefore we have the following list of relations:

- (1) $x_1^{-1}x_2x_1 = x_{n+4}x_{n+3} \cdots x_6x_2x_6^{-1} \cdots x_{n+3}^{-1}x_{n+4}^{-1}$
- (2) $x_4 = x_2x_{n+5}x_5x_{n+5}^{-1}x_2^{-1}$

- (3) $x_1^{-1}x_2^{-1}x_1x_4x_1^{-1}x_2x_1 = x_5^{-1} \cdot x_{n+6}^{-1} \cdots x_{n+m+4}^{-1} \cdot x_5 \cdot x_{n+m+4} \cdots x_{n+6} \cdot x_5$
- (4) $(x_2x_i)^2 = (x_ix_2)^2$, where $i = 1, 4$
- (5) $(x_1^{-1}x_2x_1x_4)^2 = (x_4x_1^{-1}x_2x_1)^2$
- (6) $(x_2x_i)^2 = (x_ix_2)^2$, where $6 \leq i \leq n+4$
- (7) $(x_5x_i)^2 = (x_ix_5)^2$, where $n+5 \leq i \leq n+m+4$
- (8) $[x_2x_1x_2^{-1}, x_i] = e$, where $5 \leq i \leq n+m+4$
- (9) $[x_i, x_j] = e$, where $6 \leq i \leq n+4$ and $j = 4, 5, n+5, \dots, n+m+4$
- (10) $[x_2^{-1}x_ix_2, x_j] = e$, where $6 \leq i < j \leq n+4$
- (11) $[x_4x_2x_4^{-1}, x_{n+5}] = e$
- (12) $[x_2, x_{n+5}] = e$
- (13) $[x_5x_4x_2x_4^{-1}x_5^{-1}, x_{n+6} \cdots x_{i-2}^{-1}x_{i-1}^{-1}x_ix_{i-1}x_{i-2} \cdots x_{n+6}] = e$, where $n+6 \leq i \leq n+m+4$
- (14) $[x_5x_2x_5^{-1}, x_i] = e$, where $n+6 \leq i \leq n+m+4$
- (15) $[x_{n+5}, x_i] = e$, where $n+6 \leq i \leq n+m+4$
- (16) $[x_5^{-1}x_ix_5, x_j] = e$, where $n+6 \leq i < j \leq n+m+4$
- (17) $x_{n+m+4}x_{n+m+3} \cdots x_5x_4x_2^2x_1 = e$.

By Relation (2), we can omit x_4 and replace it everywhere by

$$x_2x_{n+5}x_5x_{n+5}^{-1}x_2^{-1}.$$

We start with Relations (4). Take $(x_2x_4)^2 = (x_4x_2)^2$. This relation is rewritten as:

$$\begin{aligned} & x_2 \cdot x_2x_{n+5}x_5x_{n+5}^{-1}x_2^{-1} \cdot x_2 \cdot x_2x_{n+5}x_5x_{n+5}^{-1}x_2^{-1} = \\ & = x_2x_{n+5}x_5x_{n+5}^{-1}x_2^{-1} \cdot x_2 \cdot x_2x_{n+5}x_5x_{n+5}^{-1}x_2^{-1} \cdot x_2. \end{aligned}$$

By Relation (12) we easily get: $(x_2x_5)^2 = (x_5x_2)^2$.

Now we consider Relation (11):

$$\begin{aligned} e &= [x_4x_2x_4^{-1}, x_{n+5}] = \\ &= [x_2x_{n+5}x_5x_{n+5}^{-1}x_2^{-1} \cdot x_2 \cdot x_2x_{n+5}x_5x_{n+5}^{-1}x_2^{-1}, x_{n+5}] = \\ &= [x_5x_2x_5^{-1}, x_{n+5}]. \end{aligned}$$

By the above resulting relations, we can simplify also Relations (13):

$$\begin{aligned} e &= [x_5 \cdot x_2x_{n+5}x_5x_{n+5}^{-1}x_2^{-1} \cdot x_2 \cdot x_2x_{n+5}x_5x_{n+5}^{-1}x_2^{-1} \cdot x_5^{-1}, x_{n+6} \cdots x_{i-1}^{-1}x_ix_{i-1} \cdots x_{n+6}] = \\ &= [x_2, x_{n+6} \cdots x_{i-1}^{-1}x_ix_{i-1} \cdots x_{n+6}], \end{aligned}$$

for $n+6 \leq i \leq n+m+4$. For $i = n+6$, we get $[x_2, x_{n+6}] = e$. For $i = n+7$, we get $[x_2, x_{n+6}^{-1}x_{n+7}x_{n+6}] = e$, and by the previous relation, we get $[x_2, x_{n+7}] = e$. In the same manner, we get $[x_2, x_i] = e$, for $n+6 \leq i \leq n+m+4$.

From Relations (9), we have $[x_i, x_4] = e$, where $6 \leq i \leq n+4$. By Relation (2), $e = [x_i, x_4] = [x_i, x_2x_{n+5}x_5x_{n+5}^{-1}x_2^{-1}]$. By Relations (9) and (12), we get for $6 \leq i \leq n+4$: $[x_i, x_2x_5x_2^{-1}] = e$.

We rewrite now Relation (5), by substituting Relations (1) and (2) in it:

$$\begin{aligned} x_{n+4} \cdots x_6x_2x_6^{-1} \cdots x_{n+4}^{-1} \cdot x_2x_{n+5}x_5x_{n+5}^{-1}x_2^{-1} \cdot x_{n+4} \cdots x_6x_2x_6^{-1} \cdots x_{n+4}^{-1} \cdot x_2x_{n+5}x_5x_{n+5}^{-1}x_2^{-1} = \\ = x_2x_{n+5}x_5x_{n+5}^{-1}x_2^{-1} \cdot x_{n+4} \cdots x_6x_2x_6^{-1} \cdots x_{n+4}^{-1} \cdot x_2x_{n+5}x_5x_{n+5}^{-1}x_2^{-1} \cdot x_{n+4} \cdots x_6x_2x_6^{-1} \cdots x_{n+4}^{-1}. \end{aligned}$$

Since $[x_{n+5}, x_i] = e$, for $i = 2, 6, \dots, n+4$ (by Relations (9) and (12)),

$$\begin{aligned} x_{n+4} \cdots x_6x_2x_6^{-1} \cdots x_{n+4}^{-1}x_2x_5x_2^{-1} \cdot x_{n+4} \cdots x_6x_2x_6^{-1} \cdots x_{n+4}^{-1}x_2x_5x_2^{-1} = \\ = x_2x_5x_2^{-1}x_{n+4} \cdots x_6x_2x_6^{-1} \cdots x_{n+4}^{-1} \cdot x_2x_5x_2^{-1}x_{n+4} \cdots x_6x_2x_6^{-1} \cdots x_{n+4}^{-1}. \end{aligned}$$

Now, since we proved above that $[x_i, x_2x_5x_2^{-1}] = e$, for $6 \leq i \leq n+4$, we get $x_2x_5x_2x_5 = x_5x_2x_5x_2$, which is a consequence from Relations (4) (see above).

Therefore, we have:

Generators: $\{x_1, x_2, x_5, x_6, x_7, \dots, x_{n+4}, x_{n+5}, \dots, x_{n+m+4}\}$.

Relations:

- (1) $x_1^{-1}x_2x_1 = x_{n+4}x_{n+3} \cdots x_6x_2x_6^{-1} \cdots x_{n+3}^{-1}x_{n+4}^{-1}$
- (2) $x_1^{-1}x_2^{-1}x_1 \cdot x_2x_{n+5}x_5x_{n+5}^{-1}x_2^{-1} \cdot x_1^{-1}x_2x_1 = x_5^{-1} \cdot x_{n+6}^{-1} \cdots x_{n+m+4}^{-1} \cdot x_5 \cdot x_{n+m+4} \cdots x_{n+6} \cdot x_5$
- (3) $(x_2x_i)^2 = (x_ix_2)^2$, where $i = 1, 5, 6, \dots, n+4$
- (4) $(x_5x_i)^2 = (x_ix_5)^2$, where $n+5 \leq i \leq n+m+4$
- (5) $[x_2x_1x_2^{-1}, x_i] = e$, where $5 \leq i \leq n+m+4$
- (6) $[x_i, x_j] = e$, where $6 \leq i \leq n+4$ and $j = 5, n+5, \dots, n+m+4$
- (7) $[x_i, x_2x_5x_2^{-1}] = e$, where $6 \leq i \leq n+4$
- (8) $[x_2^{-1}x_ix_2, x_j] = e$, where $6 \leq i < j \leq n+4$
- (9) $[x_2, x_i] = e$, where $n+6 \leq i \leq n+m+4$
- (10) $[x_5x_2x_5^{-1}, x_i] = e$, where $n+5 \leq i \leq n+m+4$
- (11) $[x_{n+5}, x_i] = e$, where $i = 2, n+6, \dots, n+m+4$
- (12) $[x_5^{-1}x_ix_5, x_j] = e$, where $n+6 \leq i < j \leq n+m+4$
- (13) $x_{n+m+4}x_{n+m+3} \cdots x_5x_2x_{n+5}x_5x_{n+5}^{-1}x_2x_1 = e$.

Now we use the projective relation (Relation (13)) in order to omit the generator x_1 .

We rewrite Relation (1), by substituting x_1 :

$$\begin{aligned} x_{n+m+4} \cdots x_5 x_2 x_{n+5} x_5 x_{n+5}^{-1} x_2 x_2^{-1} x_{n+5} x_5^{-1} x_{n+5}^{-1} x_2^{-1} x_5^{-1} \cdots x_{n+m+4}^{-1} &= \\ &= x_{n+4} \cdots x_6 x_2 x_6^{-1} \cdots x_{n+4}^{-1}. \end{aligned}$$

By Relations (6) and (11), we get:

$$x_{n+m+4} \cdots x_{n+5} x_5 x_2 x_{n+5} x_5 x_2^{-1} x_{n+5}^{-1} x_2^{-1} x_5^{-1} x_{n+5}^{-1} \cdots x_{n+m+4}^{-1} = x_2.$$

By Relations (9) and (10), we have:

$$x_{n+5} x_5 x_2 x_5 x_2^{-1} x_5^{-1} x_{n+5}^{-1} = x_2.$$

Now, by Relations (3),

$$x_{n+5} x_2 x_{n+5}^{-1} = x_2,$$

which is known already in (11).

Now we simplify $(x_1 x_2)^2 = (x_2 x_1)^2$, which appears in Relations (3).

By substituting x_1 and some immediate cancellations,

$$\begin{aligned} x_{n+4} \cdots x_6 x_5 x_2 x_{n+5} x_5 x_{n+m+4} \cdots x_{n+6} x_{n+4} \cdots x_6 &= \\ &= x_2^{-1} x_{n+4} \cdots x_6 x_5 x_2 x_{n+5} x_5 x_{n+m+4} \cdots x_{n+6} x_{n+4} \cdots x_6 x_2. \end{aligned}$$

By Relations (6) and (9),

$$x_{n+4} \cdots x_6 x_5 x_2 x_{n+5} x_5 x_{n+4} \cdots x_6 = x_2^{-1} x_{n+4} \cdots x_6 x_5 x_2 x_{n+5} x_5 x_{n+4} \cdots x_6 x_2.$$

Since $[x_i, x_5] = e$ for $6 \leq i \leq n+4$ (by Relations (6)), we get

$$x_{n+4} \cdots x_6 x_5 x_2 x_{n+5} x_{n+4} \cdots x_6 x_5 = x_2^{-1} x_{n+4} \cdots x_6 x_5 x_2 x_{n+5} x_{n+4} \cdots x_6 x_5 x_2,$$

which is:

$$x_{n+4} \cdots x_6 x_5 x_2 x_{n+5} x_{n+4} \cdots x_6 x_5 x_2^{-1} x_5^{-1} = x_2^{-1} x_{n+4} \cdots x_6 x_5 x_2 x_{n+5} x_{n+4} \cdots x_6.$$

By Relations (6) and (10),

$$x_{n+4} \cdots x_6 x_5 x_2 x_{n+4} \cdots x_6 x_5 x_2^{-1} x_5^{-1} = x_2^{-1} x_{n+4} \cdots x_6 x_5 x_2 x_{n+4} \cdots x_6.$$

Now we add $x_2 x_2^{-1} = e$ in two locations:

$$x_2 x_{n+4} \cdots x_6 (x_2 x_2^{-1}) x_5 x_2 x_{n+4} \cdots x_6 x_5 x_2^{-1} x_5^{-1} = x_{n+4} \cdots x_6 (x_2 x_2^{-1}) x_5 x_2 x_{n+4} \cdots x_6.$$

By Relations (7),

$$x_2 x_{n+4} \cdots x_6 x_2 x_{n+4} \cdots x_6 x_2^{-1} x_5 x_2 x_5 x_2^{-1} x_5^{-1} = x_{n+4} \cdots x_6 x_2 x_{n+4} \cdots x_6 x_2^{-1} x_5 x_2.$$

Since $(x_2x_5)^2 = (x_5x_2)^2$ (see Relations (3)), we get:

$$x_2x_{n+4} \cdots x_6x_2x_{n+4} \cdots x_6 = x_{n+4} \cdots x_6x_2x_{n+4} \cdots x_6x_2.$$

Now we add again $x_2x_2^{-1} = e$ as follows:

$$\begin{aligned} x_2(x_2x_2^{-1})x_{n+4}(x_2x_2^{-1}) \cdots (x_2x_2^{-1})x_7(x_2x_2^{-1})x_6x_2x_{n+4} \cdots x_6 &= \\ &= x_{n+4}(x_2x_2^{-1}) \cdots (x_2x_2^{-1})x_6x_2x_{n+4} \cdots x_6x_2. \end{aligned}$$

We can rewrite the relation as follows:

$$\begin{aligned} x_2(x_2^{-1}x_{n+4}x_2)x_2^{-1} \cdots x_2(x_2^{-1}x_7x_2)(x_2^{-1}x_6x_2)x_{n+4} \cdots x_6 &= \\ &= (x_2^{-1}x_{n+4}x_2)x_2^{-1} \cdots x_2(x_2^{-1}x_7x_2)(x_2^{-1}x_6x_2)x_{n+4} \cdots x_6x_2, \end{aligned}$$

and using Relations (8), we get:

$$\begin{aligned} x_2(x_2^{-1}x_{n+4}x_2)x_{n+4} \cdots (x_2^{-1}x_7x_2)x_7(x_2^{-1}x_6x_2)x_6 &= \\ &= (x_2^{-1}x_{n+4}x_2)x_{n+4} \cdots (x_2^{-1}x_7x_2)x_7(x_2^{-1}x_6x_2)x_6x_2. \end{aligned}$$

Since by Relations (3) we have $(x_2x_i)^2 = (x_ix_2)^2$ for $6 \leq i \leq n+4$, this relation is redundant.

Relation (2) can be simplified by substituting Relation (1) and using Relations (6) and (11):

$$\begin{aligned} x_{n+5} \cdot x_{n+4} \cdots x_6 \cdot x_2^{-1} \cdot x_6^{-1} \cdots x_{n+4}^{-1} \cdot x_2x_5x_2^{-1} \cdot x_{n+4} \cdots x_6 \cdot x_2 \cdot x_6^{-1} \cdots x_{n+4}^{-1} \cdot x_{n+5} &= \\ &= x_5^{-1} \cdot x_{n+6}^{-1} \cdots x_{n+m+4}^{-1} \cdot x_5 \cdot x_{n+m+4} \cdots x_{n+6} \cdot x_5. \end{aligned}$$

By Relations (6) and (7),

$$x_5x_{n+5}x_5x_{n+5}^{-1}x_5^{-1} = x_{n+6}^{-1} \cdots x_{n+m+4}^{-1} \cdot x_5 \cdot x_{n+m+4} \cdots x_{n+6}.$$

Using again $(x_5x_{n+5})^2 = (x_{n+5}x_5)^2$, we get:

$$x_{n+5}^{-1}x_5x_{n+5} = x_{n+6}^{-1} \cdots x_{n+m+4}^{-1} \cdot x_5 \cdot x_{n+m+4} \cdots x_{n+6},$$

which can be rewritten as:

$$(*) \quad x_{n+m+4} \cdots x_{n+6} \cdot x_5 \cdot x_{n+5} = x_{n+5} \cdot x_5 \cdot x_{n+m+4} \cdots x_{n+6}.$$

We will see now that Relations (5) are all redundant. We split it into four subcases, according to the value of i :

- If we substitute $i = 5$ in Relations (5), we have $[x_1, x_2^{-1}x_5x_2] = e$, which can be simplified to:

$$x_{n+m+4} \cdots x_5x_2x_{n+5}x_5x_{n+5}^{-1}x_5 = x_2^{-1}x_5x_2x_{n+m+4} \cdots x_5x_2x_{n+5}x_5x_{n+5}^{-1}.$$

By adding $x_5^{-1}x_5 = e$ and using $(x_5x_{n+5})^2 = (x_{n+5}x_5)^2$, we get:

$$\begin{aligned} & x_{n+m+4} \cdots x_{n+6}x_{n+5}x_{n+4} \cdots x_6x_5x_2x_5^{-1}x_{n+5}^{-1}x_5x_{n+5}x_5 = \\ & = x_2^{-1}x_5x_2x_{n+m+4} \cdots x_{n+6}x_{n+5}x_{n+4} \cdots x_6x_5x_2x_5^{-1}x_{n+5}^{-1}x_5x_{n+5}. \end{aligned}$$

By Relations (6), (7) and (10),

$$x_{n+m+4} \cdots x_{n+6}x_5x_2x_{n+5}x_5x_{n+5}^{-1}x_5^{-1} = x_2^{-1}x_5x_2x_{n+m+4} \cdots x_{n+6}x_5x_2x_5^{-1}.$$

By Relations (10), (11) and $(x_2x_5)^2 = (x_5x_2)^2$:

$$x_{n+m+4} \cdots x_{n+6}x_5x_{n+5}x_5x_{n+5}^{-1}x_5^{-1}x_2 = x_5x_2x_{n+m+4} \cdots x_{n+6}.$$

By Relations (9) and $(x_5x_{n+5})^2 = (x_{n+5}x_5)^2$,

$$x_{n+5}^{-1}x_5x_{n+5} = x_{n+6}^{-1} \cdots x_{n+m+4}^{-1}x_5x_{n+m+4} \cdots x_{n+6},$$

which is known already (by (*)).

- For $6 \leq i \leq n+4$, we have $[x_1, x_2^{-1}x_ix_2] = e$. By some cancellations, we have:

$$x_{n+m+4} \cdots x_5x_2x_{n+5}x_5x_{n+5}^{-1}x_i = x_2^{-1}x_ix_2x_{n+m+4} \cdots x_5x_2x_{n+5}x_5x_{n+5}^{-1}.$$

By Relations (6) and (9), $x_{n+4} \cdots x_6x_5x_2x_i = x_2^{-1}x_ix_2x_{n+4} \cdots x_6x_5x_2$.

In this relation we substitute first $i = 6$ to get (by Relations (8)):

$$x_6x_5x_2x_6 = x_2^{-1}x_6x_2x_6x_5x_2,$$

which is $[x_6, x_2^{-1}x_5x_2] = e$. This relation appears already in Relations (7). Now we substitute $i = 7$ to get (again by Relations (8)):

$$x_7x_6x_5x_2x_7 = x_2^{-1}x_7x_2x_7x_6x_5x_2,$$

which is $x_6x_5x_2x_7 = x_2x_7x_2^{-1}x_6x_5x_2$. The addition of $x_2x_2^{-1} = e$ in two locations, enables us to translate

$$x_2^{-1}x_6(x_2x_2^{-1})x_5x_2x_7 = x_7x_2^{-1}x_6(x_2x_2^{-1})x_5x_2$$

to $(x_2^{-1}x_6x_2)(x_2^{-1}x_5x_2)x_7 = x_7(x_2^{-1}x_6x_2)(x_2^{-1}x_5x_2)$, which is of course redundant by Relations (8). As for the other indices, all relations are redundant in a similar way.

- If $i = n + 5$, we have by Relations (11): $[x_1, x_{n+5}] = e$. By some immediate cancellations, we get:

$$x_{n+m+4}x_{n+m+3} \cdots x_5x_2 = x_{n+5}x_{n+m+4}x_{n+m+3} \cdots x_5x_2x_{n+5}x_5x_{n+5}^{-1}x_5^{-1}x_{n+5}^{-1}.$$

By Relations (6) and (11), we have $x_5x_2 = x_{n+5}x_5x_2x_{n+5}x_5x_{n+5}^{-1}x_5^{-1}x_{n+5}^{-1}$.

By Relations (4) and (10), the relation is redundant.

- For $n+6 \leq i \leq n+m+4$, we have by Relations (9): $[x_1, x_i] = e$. Substituting Relation (13), we have:

$$x_{n+m+4} \cdots x_5x_2x_{n+5}x_5x_{n+5}^{-1}x_2x_i = x_ix_{n+m+4} \cdots x_5x_2x_{n+5}x_5x_{n+5}^{-1}x_2,$$

which is (by Relations (6), (9) and (11)):

$$x_{n+m+4} \cdots x_{n+6}x_5x_2x_{n+5}x_5x_i = x_ix_{n+m+4} \cdots x_{n+6}x_5x_2x_{n+5}x_5.$$

We add $x_5^{-1}x_5 = e$ in two locations as follows:

$$x_{n+m+4} \cdots x_{n+6}x_5x_2(x_5^{-1}x_5)x_{n+5}x_5x_i = x_ix_{n+m+4} \cdots x_{n+6}x_5x_2(x_5^{-1}x_5)x_{n+5}x_5.$$

This enables us to use Relations (10) and to cancel $x_5x_2x_5^{-1}$:

$$x_{n+m+4} \cdots x_{n+6}x_5x_{n+5}x_5x_i = x_ix_{n+m+4} \cdots x_{n+6}x_5x_{n+5}x_5.$$

By Relation (*), we can rewrite it as:

$$x_{n+5}x_5x_{n+m+4} \cdots x_{n+6}x_5x_i = x_ix_{n+5}x_5x_{n+m+4} \cdots x_{n+6}x_5.$$

By relations (11), we get $[x_5^{-1}x_ix_5, x_{n+m+4} \cdots x_{n+6}x_5^2] = e$.

Now we substitute $i = n + 6$:

$$x_5^{-1}x_{n+6}x_5x_{n+m+4} \cdots x_{n+6}x_5^2 = x_{n+m+4} \cdots x_{n+6}x_5^2x_5^{-1}x_{n+6}x_5.$$

By Relations (12), we get $x_{n+6}x_5x_{n+6}x_5 = x_5x_{n+6}x_5x_{n+6}$, which is known already (by Relations (4)).

Now we substitute $i = n + 7$:

$$x_5^{-1}x_{n+7}x_5x_{n+m+4} \cdots x_{n+6}x_5 = x_{n+m+4} \cdots x_{n+6}x_5x_{n+7},$$

Again, by Relations (12), we get:

$$x_5^{-1}x_{n+7}x_5x_{n+7}x_{n+6}x_5 = x_{n+7}x_{n+6}x_5x_{n+7},$$

and by $(x_5x_{n+7})^2 = (x_{n+7}x_5)^2$ (Relations (4)), we get:

$$x_5x_{n+7}x_5^{-1}x_{n+6}x_5 = x_{n+6}x_5x_{n+7}.$$

Now, since $[x_{n+7}, x_5^{-1}x_{n+6}x_5] = e$ (Relations (12)), this relation is redundant.

In a similar way, the relations $[x_1, x_i] = e$ for $n+6 \leq i \leq n+m+4$, are redundant.

Therefore the resulting presentation is as follows:

Generators: $\{x_2, x_5, x_6, x_7, \dots, x_{n+4}, x_{n+4}, x_{n+5}, \dots, x_{n+m+4}\}$.

Relations:

- (1) $x_{n+5}^{-1}x_5x_{n+5} = x_{n+6}^{-1} \cdots x_{n+m+4}^{-1} \cdot x_5 \cdot x_{n+m+4} \cdots x_{n+6}$
- (2) $(x_2x_i)^2 = (x_ix_2)^2$, where $i = 5, 6, \dots, n+4$
- (3) $(x_5x_i)^2 = (x_ix_5)^2$, where $n+5 \leq i \leq n+m+4$
- (4) $[x_i, x_j] = e$, where $6 \leq i \leq n+4$ and $j = 5, n+5, \dots, n+m+4$
- (5) $[x_i, x_2x_5x_2^{-1}] = e$, where $6 \leq i \leq n+4$
- (6) $[x_2^{-1}x_ix_2, x_j] = e$, where $6 \leq i < j \leq n+4$
- (7) $[x_2, x_i] = e$, where $n+6 \leq i \leq n+m+4$
- (8) $[x_5x_2x_5^{-1}, x_i] = e$, where $n+5 \leq i \leq n+m+4$
- (9) $[x_{n+5}, x_i] = e$, where $i = 2, n+6, \dots, n+m+4$
- (10) $[x_5^{-1}x_ix_5, x_j] = e$, where $n+6 \leq i < j \leq n+m+4$,

as stated in Theorem 1.1. ■

As we have seen in Corollary 3.10, the projective fundamental groups of this section are big.

Remark 4.6. *Note that if we substitute $m = 0$ in the presentation of Theorem 1.1, we get exactly the presentation of $T_{n,0}$ given in Theorem 3.9.*

5. ACKNOWLEDGEMENTS

We would like to thank Mutsuo Oka for correcting a wrong assumption in the first version of this paper.

REFERENCES

- [1] Amram, M., Garber, D. and Teicher, M., *Fundamental groups of tangent conic-line arrangements with singularities up to order 6*, Math. Z. **256**, 837–870 (2007).
- [2] Amram, M. and Ogata, S., *Degenerations and fundamental groups related to some special toric varieties*, Michigan Math. J. **54**, 587–610 (2006).
- [3] Amram, M. and Teicher, M., *Braid monodromy of special curves*, J. Knot Theory & Ramifications **10**(2), 171–212 (2001).
- [4] Amram, M. and Teicher, M., *Fundamental groups of some special quadric arrangements*, Rev. Mat. Complut. **19**(2), 259–276 (2006).
- [5] Amram, M., Teicher, M. and Uludag, M., *Fundamental groups of some quadric-line arrangements*, Topology Appl. **130**, 159–173 (2003).
- [6] Artal-Bartolo, E., *Sur les couples de Zariski*, J. Alg. Geom. **3**(2), 223–247 (1994).
- [7] Artal-Bartolo, E. and Carmona-Rubel, J., *Zariski pairs, fundamental groups and Alexander polynomials*, J. Math. Soc. Japan **50**(3), 521–543 (1998).
- [8] Chisini, O., *Sulla identità birazionale delle funzioni algebriche di due variabili dotate di una medesima curva di diramazione*, Rend. Ist. Lombardo **77**, 339–356 (1944).
- [9] Degtyarev, A. I., *Quintics in $\mathbb{CP}^2$ with non-abelian fundamental group*, Algebra i Analiz **11**(5), 130–151 (1999) [Russian]; *Translation in*: St. Petersburg Math. J. **11**(5), 809–826 (2000).
- [10] Garber, D., *Plane curves and their fundamental groups: Generalizations of Uludag’s Construction*, Alg. Geom. Topo. **3**, 593–622 (2003).
- [11] Garber, D. and Teicher, M., *The fundamental group’s structure of the complement of some configurations of real line arrangements*, Complex Analysis and Algebraic Geometry, edited by T. Peternell and F.-O. Schreyer, de Gruyter, 173–223 (2000).
- [12] Hirzebruch, F., *Some examples of algebraic surfaces*, Contemp. Math. **9**, 55–71 (1982).
- [13] Kulikov, V. S., *On Chisini’s conjecture*, Izv. Ross. Akad. Nauk Ser. Mat. **63**:6, 83–116 (1999) [Russian]; *Translation in* Izv. Math. **63**, 1139–1170 (1999).
- [14] Kulikov, V. S. and Teicher, M., *Braid monodromy factorizations and diffeomorphism types*, Izv. Ross. Akad. Nauk Ser. Mat. **64**(2), 89–120 (2000) [Russian]; *Translation in* Izv. Math. **64**(2), 311–341 (2000).
- [15] Milnor, J., *Morse Theory*, Ann. Math. Stud. **51**, Princeton University Press, Princeton, NJ, 1963.
- [16] Moishezon, B., *Stable branch curves and braid monodromies*, Lect. Notes in Math. **862**, 107–192 (1981).

- [17] Moishezon, B. and Teicher, M., *Braid group techniques in complex geometry I, Line arrangements in $\mathbb{CP}^2$* , Contemp. Math. **78**, 425–555 (1988).
- [18] Moishezon, B. and Teicher, M., *Braid group techniques in complex geometry II, From arrangements of lines and conics to cuspidal curves*, Algebraic Geometry, Lect. Notes in Math. **1479**, 131–180 (1990).
- [19] Moishezon, B. and Teicher, M., *Braid group techniques in complex geometry III, Projective degeneration of V_3* , Contemp. Math. **162**, 313–332 (1994).
- [20] Moishezon, B. and Teicher, M., *Braid group techniques in complex geometry IV, Braid monodromy of the branch curve S_3 of $V_3 \rightarrow \mathbb{CP}^2$ and application to $\pi_1(\mathbb{CP}^2 - S_3, *)$* , Contemp. Math. **162**, 332–358 (1994).
- [21] Oka, M., *Two transforms of plane curves and their fundamental groups*, J. Math. Sci. Univ. Tokyo **3**(2), 399–443 (1996).
- [22] Shimada, I., *A note on Zariski pairs*, Compositio Math. **104**(2), 125–133 (1996).
- [23] Tokunaga, H., *Some examples of Zariski pairs arising from certain elliptic K3 surfaces*, Math. Z. **227**(3), 465–477 (1998).
- [24] Uludağ, A. M., *More Zariski pairs and finite fundamental group of curve complements*, Manu. Math. **106**(3), 271–277 (2001).
- [25] van Kampen, E. R., *On the fundamental group of an algebraic curve*, Amer. J. Math. **55**, 255–260 (1933).
- [26] Zariski, O., *On the problem of existence of algebraic functions of two variables possessing a given branch curve*, Amer. J. Math. **51**, 305–328 (1929).
- [27] Zariski, O., *The topological discriminant group of a Riemann surface of genus p* , Amer. J. Math. **59**, 335–358 (1937).

MEIRAV AMRAM AND DAVID GARBER, DEPARTMENT OF APPLIED MATHEMATICS, FACULTY OF SCIENCES, HOLON INSTITUTE OF TECHNOLOGY, 52 GOLOMB ST., PO BOX 305, 58102 HOLON, ISRAEL

E-mail address: ameirav@math.huji.ac.il, garber@hit.ac.il

MINA TEICHER, DEPARTMENT OF MATHEMATICS, BAR-ILAN UNIVERSITY, 52900 RAMAT-GAN, ISRAEL

E-mail address: teicher@macs.biu.ac.il